

**This Page Is Inserted by IFW Operations  
and is not a part of the Official Record**

## **BEST AVAILABLE IMAGES**

**Defective images within this document are accurate representations of the original documents submitted by the applicant.**

**Defects in the images may include (but are not limited to):**

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/00, C12N 1/15, 15/63, C12P 21/02</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/25735</b>
		<b>(43) International Publication Date:</b> 27 May 1999 (27.05.99)
<b>(21) International Application Number:</b> PCT/US98/24975 <b>(22) International Filing Date:</b> 19 November 1998 (19.11.98)  <b>(30) Priority Data:</b> 60/066,129 19 November 1997 (19.11.97) US 60/066,308 21 November 1997 (21.11.97) US 60/066,462 24 November 1997 (24.11.97) US  <b>(71) Applicant:</b> MICROBIA, INC. [US/US]; 840 Memorial Drive, Cambridge, MA 02139 (US).  <b>(72) Inventors:</b> HECHT, Peter; 17 Duffield Road, Newton, MA 02166 (US). MADDEN, Kevin; 174 Newbury Street #6, Boston, MA 02116 (US). FINK, Gerald, R.; 40 Aston Road, Chestnut Hill, MA 02167 (US).  <b>(74) Agent:</b> CLARK, Paul, T.; Clark & Elbing LLP, 176 Federal Street, Boston, MA 02110-2214 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>

**(54) Title:** CHIMERIC PRE-ACTIVATED TRANSCRIPTION FACTORS**(57) Abstract**

Disclosed herein is a chimeric protein comprising a pre-activated transcription factor and a strong transcription activation domain for regulating fungal gene expression, and reagents and methods for constructing and using said protein.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## CHIMERIC PRE-ACTIVATED TRANSCRIPTION FACTORS

### Background of the Invention

5 Fungal species are the commercial source of many medicinally useful products, such as antibiotics (e.g., beta-lactam antibiotics such as penicillin, cephalosporin, and their derivatives), anti-hypercholesterolemic agents (e.g., lovastatin and compactin), immunosuppressives (e.g., cyclosporin), and antifungal drugs (e.g., pneumocandin and echinocandin). All  
10 of these drugs are fungal secondary metabolites, small secreted molecules that fungi utilize against competitors in their microbial environment. Fungi also produce commercially important enzymes (e.g., cellulases, proteases, and lipases) and other products (e.g., citric acid, gibberellic acid, natural pigments, and flavorings).

15 The production of secondary metabolites, enzymes, and other products is regulated by coordinated gene expression. For example, the production of penicillin is limited by the activity of two enzymes, encoded by the *ipnA* and *acvA* genes. PacC, a zinc-finger transcription factor, binds to sequences upstream of these two genes. Moreover, increased activity of PacC  
20 leads to both increased enzyme activity and penicillin production.

Our understanding of transcriptional regulation of secondary metabolite production, as exemplified above, has increased greatly over the past decade. To date, however, the use of genetically-engineered transcription factors has not been applied to increase production of commercially-important  
25 fungal products. In contrast, methods to increase production of penicillin currently rely upon mutagenesis and selection for mutants which display increased secondary metabolite production.

-2-

Summary of the Invention

The invention provides a means to increase the production of secondary metabolites in fungi by genetic manipulation of the fungal organism itself. The ability to increase fungal secondary metabolite production has at least two important applications. First, it will allow increased production of existing secondary metabolites which are useful in clinical and experimental settings. Second, increasing production of secondary metabolites will facilitate identification of new compounds in fungi that otherwise make undetectable levels of these compounds in the laboratory.

Accordingly, in one aspect, the invention features a two-part chimeric transcription factor including (i) a pre-activated transcription factor functional in a fungal strain, and (ii) a transcription activation domain that is different from the transcription activation domain naturally associated with the transcription factor. In a preferred embodiment, the transcriptional activity of the chimeric transcription factor is greater than the transcriptional activity naturally associated with the pre-activated transcription factor. In another preferred embodiment, the pre-activated transcription factor is pre-activated by truncation. In a related preferred embodiment, the pre-activated transcription factor includes a substitution of a serine or threonine residue with an alanine, aspartic acid, or glutamic acid residue, wherein the substitution pre-activates the transcription factor (e.g., by mimicking or otherwise altering phosphorylation). In another preferred embodiment, the transcription factor is a member of the PacC family (defined below) and can be pre-activated. In a related preferred embodiment, the pre-activated transcription factor contains portions of the amino acid sequence shown in Fig. 1 (SEQ ID NOs: 1-6).

In another aspect, the invention features a vector including DNA encoding a chimeric transcription factor including (i) a pre-activated

-3-

transcription factor functional in a fungal strain, and (ii) a transcription activation domain that is different from the transcription activation domain naturally associated with the transcription factor. The DNA is operably linked to a promoter capable of directing and regulating expression of the chimeric  
5 transcription factor in a fungal strain.

The transcription factor encoded within the vector described above is expressed in a fungal cell, such as a filamentous fungal cell, which produces the secondary metabolite of interest and in which expression of the transcription factor increases the production of the secondary metabolite by the  
10 cell. The secondary metabolite can be non-proteinaceous or it can be a protein or peptide.

In another aspect, the invention features a method of producing a secondary metabolite of interest, including the steps of (i) introducing into a fungal cell, such as a filamentous fungal cell, a vector including a promoter  
15 capable of controlling gene expression in the fungal cell, and a nucleic acid encoding a two-part transcription factor including a DNA-binding domain and a transcription activation domain; and (ii) culturing the fungal cell under secondary metabolite-producing conditions. In a preferred embodiment, the transcription activation domain is different from the transcription activation  
20 domain naturally associated with the DNA-binding domain. In other preferred embodiments, the transcription factor is a pre-activated transcription factor (pre-activated by substitution of a serine or threonine residue with an alanine, aspartic acid, or glutamic acid residue, or pre-activated by truncation). In other preferred embodiments, the DNA binding domain of the transcription factor is  
25 from a fungal transcriptional activator or from a fungal transcriptional repressor.

By "pre-activated transcription factor" is meant a transcription factor

-4-

or fragment thereof that, compared to the precursor molecule, is capable of 1) increased binding, either direct or indirect, to a specific DNA sequence located in a gene regulatory region (e.g., a promoter), or 2) increased transcription activating properties. Pre-activated transcription factors may be able to activate transcription from promoters, but this is not necessarily the case. For example, a transcription factor DNA-binding domain with binding properties but no transactivation activity is considered to be a pre-activated transcription factor. "Pre-activation by truncation" or "pre-activated by truncation" means that removal of a portion of the protein leads to pre-activation. This occurs *in vivo* through proteolytic cleavage. In the invention, pre-activation by truncation is achieved with the use of DNA that encodes a pre-activated form of the protein, excluding portions of the protein that would be proteolytically cleaved *in vivo*.

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 50%, preferably 85%, more preferably 90%, and most preferably 95% identity to a reference amino acid or nucleic acid sequence. For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides.

By "promoter" is meant a sequence sufficient to direct and/or regulate transcription. Also included in the invention are those elements which are sufficient to render promoter-dependent gene expression controllable for cell type-specific, tissue-specific, temporal-specific, or inducible by external signals or agents; such elements may be located in the 5' or 3' or intron



-5-

sequence regions of the native gene.

By "operably linked" is meant that a gene and one or more regulatory sequences are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequences.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

#### Drawing

Fig. 1 is an alignment of the zinc-finger DNA-binding domain of PacC family members from *Aspergillus nidulans* (SEQ ID NO: 1), *Aspergillus niger* (SEQ ID NO: 2), *Penicillium chrysogenum* (SEQ ID NO: 3), *Yarrowia lipolytica* (SEQ ID NO: 4), *Candida albicans* (SEQ ID NO: 5), and *Saccharomyces cerevisiae* (SEQ ID NO: 6). Identity is represented by shaded regions; similarity is represented by boxed regions.

#### Detailed Description

The invention features a two-part chimeric protein including a pre-activated transcription factor and a strong transcription activation domain for regulating fungal gene expression. The protein is encoded by nucleic acids operably linked to a strong promoter in a vector which allows for expression in fungal cells. The effect of the transcription factor is to facilitate expression of a protein which itself is a desired product, or which acts as an element (e.g., an enzyme) by which a desired product is made by the host fungal cell. Each of these components is described below. Experimental examples described herein are intended to illustrate, not limit, the scope of the claimed invention.

-6-

Pre-Activated Transcription Factor

The vectors of the invention can include DNA encoding any proteinaceous transcription factor that can be provided in pre-activated form; i.e., the vector encodes the protein in a form in which it is already activated; 5 i.e., no post-translational processing is required for the protein to be active in a fungal cell to bind to regulatory DNA of the cell to facilitate gene expression.

Transcription factors regulate the level of gene expression by affecting the activity of the core transcriptional machinery at the promoter of each gene. Several mechanisms have evolved to control the activity of 10 transcription factors.

Post-translational modification is one mechanism by which transcription factors are regulated. Proteolytic cleavage is one post-translational mechanism for regulating the activity of a transcription factor (e.g., Pahl and Baeuerle, *Curr. Opin. Cell Biol.*, 1996, 8:340-347; Goodbourn 15 and King, *Biochem. Soc. Trans.*, 1997, 25:498-502; Fan and Maniatis, *Nature*, 1991, 354:395-398). The fungal PacC family of transcription factors is one class of proteins that can be activated by proteolysis. Activating mutations have been described for PacC family members (see below); these mutations truncate the encoded protein, resulting in the production of a pre-activated form 20 of the transcription factor.

Another method for pre-activating a transcription factor is to mimic the modifications which normally regulate its activity. For example, phosphorylation has been shown to positively regulate the activity of some transcription factors and negatively regulate that of others (see review by 25 Hunter and Karin, *Cell*, 1992, 70:375-387). Other forms of post-translational modifications that can increase the activity of transcription factors include acetylation (Gu and Roeder, *Cell*, 1997, 90:595-606) and alkylation (e.g.,

-7-

methylation)(Chinenov et al., *J. Biol. Chem.*, 1998, 273:6203-6209; Sakashita et al., *J Biochem (Tokyo)*, 1995, 118:1184-1191).

Dephosphorylation of particular residues can increase the activity of many transcription factors. Phosphorylation most commonly occurs on serine (Ser), threonine (Thr), and tyrosine (Tyr) residues; in some instance residues such as aspartate (Asp) and histidine (His) can be phosphorylated. The coding sequence for the phosphorylated residue can be mutated to encode an amino acid that cannot be phosphorylated and does not have a negatively charged side chain (e.g., alanine (Ala)). Ser→Ala, Thr→Ala, Tyr→Ala, and Asp→Ala substitutions are frequently used in the art to produce a pre-activated transcription factor (see, for example, Chen et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1998, 95:2349-2354; Song et al., *Mol. Cell Biol.*, 1998, 18:4994-4999; O'Reilly et al., *EMBO J.*, 1997, 16:2420-2430; Hao et al., *J. Biol. Chem.*, 1996, 271:29380-29385).

Phosphorylation can also increase the activity of a transcription factor. Mutations of Glu or Asp for Ser, Thr, or Tyr are frequently used in the art to mimic a phosphorylation event and pre-activate a transcription factor (see, for example, Hoeffler et al., *Nucleic Acids Res.*, 1994, 22:1305-12; Hao et al., *supra*). Mutations that result in a substitution of Glu for Asp, at Asp residues which can be phosphorylated, can also cause activation (Klose et al., *J. Mol. Biol.*, 1993, 232:67-78; Krems et al., *Curr. Genet.*, 1996, 29:327-34; Nohaile et al., *J. Mol. Biol.*, 1997, 273:299-316).

Other mutations can be made that mimic activating post-translational modifications. For example, the *E. coli* Ada transcription factor is activated by methylation of cysteine (Cys) residue 69. A Cys→His substitution was found to result in activation (Taketomi et al., *Mol. Gen. Genet.*, 1996, 250:523-532). This particular substitution was identified by substituting Cys 69 with each of

-8-

the other nineteen amino acids. Alternatively, in instances where no obvious substitution can be made to mimic a modification (e.g., acetylation), a random mutagenesis is performed to identify constitutively active forms of transcription factors (see, for example, Onishi et al., *Mol. Cell Biol.*, 1998, 18:3871-3879).

- 5 This technique can employ simple and rapid phenotypic or reporter selections, such as those described herein, to identify activated forms. For example, a *Saccharomyces cerevisiae* strain containing a reporter construct can be used to select for activated forms. Specifically, the *ipnA* promoter ( $P_{ipnA}$ ) from *Aspergillus nidulans* may be fused to a gene from *Saccharomyces cerevisiae*
- 10 that confers a growth advantage, such as *HIS3*, when PacC is pre-activated by a mutation. A  $P_{ipnA}$ -*HIS3* fusion has the added advantage that expression levels can be titrated by the compound 3-aminotriazole (3-AT). 3-AT is a competitive inhibitor of His3 that, when present in sufficient amounts, will inhibit the His3 expressed from  $P_{ipnA}$  and prevent this strain from growing on
- 15 SC-HIS. In this example, *pacC* coding sequence can be randomly mutagenized and vectors containing the mutated alleles are transformed into the reporter strain. Growth of a strain containing  $P_{ipnA}$ -*HIS3* only occurs on SC-HIS+3-AT plates when  $P_{ipnA}$ -*HIS3* expression is increased to overcome the competitive inhibition of His3 by 3-AT. This method provides a rapid technique for
- 20 screening for mutations which pre-activate a transcription factor.

#### The PacC Family of Transcription Factors

- One group of transcription factors useful in the invention are members of the PacC family. The PacC transcription factors regulate gene expression in response to changes in ambient pH. Members of the family have
- 25 the following characteristics: 1) They display significant (at least 35%) amino acid sequence identity to the *Aspergillus nidulans* PacC protein (Tilburn et al.,

-9-

*EMBO J.*, 1995, 14:779-790). Such proteins have been identified in *Yarrowia lipolytica* (YlRim101p; Lambert et al., *Mol. Cell. Biol.*, 1997, 17:3966-3976), *Penicillium chrysogenum* (Suarez and Penalva, *Mol. Microbiol.*, 1996, 20:529-540), *Aspergillus niger* (MacCabe et al., *Mol. Gen. Genet.*, 1996, 250:367-374),  
 5 *Saccharomyces cerevisiae* (Inv8/Rim101/Rim1; Su and Mitchell, *Nucleic Acids Res.*, 1993, 21:3789-3797), and *Candida albicans* (U.S.S.N. \_\_\_\_/\_\_\_\_)(Table 1). 2) They contain a predicted DNA-binding region that includes three zinc fingers of the Cys<sub>2</sub>His<sub>2</sub> class.

TABLE 1

10	Species of origin of PacC homolog	% identity to <i>A. nidulans</i> PacC in 107-aa	% similarity to <i>A. nidulans</i> PacC over entire length
	<i>A. Niger</i>	94	75
	<i>P. chrysogenum</i>	84	67
	<i>C. albicans</i>	61	18
15	<i>S. cerevisiae</i>	56	22
	<i>Y. lipolytica</i>	58	30

In addition, several PacC family member either have been shown to directly bind to or regulate expression of genes that contain a 5'-GCCAAG-3' or 5'-GCCAGG-3' element in upstream regulatory sequence (Tilburn et al., *supra*; Suarez  
 20 and Penalva, *supra*). Furthermore, with the exception of PacC from *P. chrysogenum*, mutations that truncate the protein have either been identified or constructed, and these mutations result in activation of gene expression by the PacC family of proteins, even at low ambient pH (Tilburn et al., *supra*; van den Hombergh et al., *Mol. Gen. Genet.*, 1996, 251:542-550; Lambert et al., *supra*; Li and Mitchell, *Genetics*, 1997,  
 25 145:63-73). Finally, in both *A. nidulans* and *S. cerevisiae*, it has been demonstrated

-10-

that specific proteolytic cleavage results in activation of signaling *in vivo* (Orejas et al., *Genes Dev.*, 1995, 9:1622-32; Li and Mitchell, *supra*).

### Transcription Activation Domains

Transcription activation domains (TADs) are discrete regions of proteins which promote gene expression by a variety of mechanisms that ultimately result in the activation of RNA polymerase. A TAD generally is defined as the minimal motif that activates transcription when fused to a DNA-binding domain (DBD) (Webster et al., *Cell*, 1988, 52:169-178; Fischer et al., *Nature*, 1988, 332:853-856; Hope et al., *Nature*, 1988, 333:635-640). The invention can employ any TAD that can transactivate expression from a fungal gene promoter when the TAD is fused to an appropriate DBD. TADs are classified based on similarities in protein sequence and/or composition properties. These classes include the acidic-rich (e.g., Gal4, Gcn4, VP16, and Jun; Webster et al., *supra*; Fischer et al., *supra*; Hope et al., *supra*; Cress and Triezenberg, *Science*, 1991, 251:87-90; Struhl, *Nature*, 1988, 332:649-650), glutamine-rich (Sp1, Oct1, and Oct2; Courey and Tjian, *Cell*, 1988, 55:887-898; Tanaka et al., *Mol. Cell Biol.*, 1994, 14:6046-6055; Tanaka and Herr, *Mol. Cell Biol.*, 1994, 14:6056-6067), and proline-rich TADs (CTF, NF-I, and EKLF; Mermod et al., *Cell*, 1989, 58:741-753; Tanese et al., *Genes Dev.*, 1991, 5:2212-2224; Chen and Bieker, *EMBO J.*, 1996, 15:5888-5896). Any of these classes of TADs may be used in the present invention. The ability of any particular TAD to transactivate from a particular promoter can be determined using simple, known selection screens.

It is also possible to artificially create either a TAD or a site-specific DBD. In one example, protein sequences which transactivate a reporter gene from a promoter of interest are selected from an expression library. In another example, protein sequences which specifically bind particular DNA sequences are selected. In each case, these sequences can then be mutated in a reiterative process to obtain either the optimal TAD sequence for the particular promoter, or the optimal DBD sequence for a particular DNA sequence. Transcription factors containing artificial elements

-11-

produced by this or any other method are useful in the invention.

In the chimeric transcription factor of the featured invention, TADs may be used alone or in combination. For example, Sp1 contains multiple glutamine-rich TADs, and these domains act synergistically to promote gene expression (Courey and Tjian, *supra*; Courey et al., *Cell*, 1989, 59:827-836). Oct-2 contains both glutamine-rich and proline-rich TADs, and both are required for maximal expression when fused to either the Oct-2 or a heterologous DBD (Tanaka et al., *supra*). Thus, the use of two or more classes of TADs in one construct may amplify the induction of expression. Furthermore, homopolymeric stretches of proline or glutamine function as TADs (Gerber et al., *Science*, 1994, 263:808-811). In one example, a strong transcription factor has been created by fusion of the Gal4 DBD to a homopolymeric glutamine stretch linked to reiterated VP16 TADs (Schwechheimer et al., *Plant Mol. Biol.*, 1998, 36:195-204).

#### Fungal Promoters

The chimeric, pre-activated transcription factor is operably linked to a strong promoter, allowing for expression of the transcription factor in a fungal cell. Expression systems utilizing a wide variety of promoters in many fungi are known, including, but not limited to, *Aspergillus nidulans* (*gpd*: Punt et al., *Gene*, 1987, 56:117-124; Hunter et al., *Curr. Genet.*, 1992, 22:377-383; Glumoff et al., *Gene*, 1989, 84:311-318. *alcA*; Fernandez-Abalos et al., *Mol. Microbiol.*, 1998, 27:121-130. *glaA*: Carrez et al., *Gene*, 1990, 94:147-154. *amdS*: Turnbull et al., *Appl. Environ. Microbiol.*, 1990, 56:2847-2852), *Aspergillus niger* (*gpd*: Punt et al., *supra*; Hunter et al., *supra*; Glumoff et al., *supra*. *glaA*: Tang et al., *Chin. J. Biotechnol.*, 1996, 12:131-136. *amdS* promoter: Turnbull et al., *supra*), *Pichia pastoris* (alcohol oxidase I promoter: Payne et al., *Gene*, 1988, 62:127-134), *Pleurotus ostreatus* (*Lentinus edodes* *ras* promoter: Yanai et al., *Biosci. Biotechnol. Biochem.*, 1996, 60:472-475), *Phytophthora infestans* (*Bremia lactucae* *Hsp70*: Judelson et al., *Mol. Plant Microbe Interact.*, 1991, 4:602-607), *Neurospora crassa* (*his3* promoter: Avalos et al., *Curr.*

-12-

*Genet.*, 1989, 16:369-372), *Yarrowia lipolytica* (*XPR2* promoter: Nicaud et al., *Curr. Genet.*, 1989, 16:253-260. *TEF*: Muller et al., *Yeast*, 1998, 14:1267-1283.), *Penicillium chrysogenum* (*phoA* promoter: Graessle et al., *Appl. Environ. Microbiol.*, 1997, 63:753-756), *Rhizopus deleamar* (*pyr4* promoter: Horiuchi et al., *Curr. Genet.*, 5 1995, 27:472-478), *Gliocladium virens* (*prom1*: Dave et al., *Appl. Microbiol. Biotechnol.*, 1994, 41:352-358), and *Cochliobolus heterostrophus* (Monke and Shafer, *Mol. Gen. Genet.*, 1993, 241:73-80).

There are also simple techniques for isolating promoters in organisms with relatively unstudied genetics. One of these is a system based on selection of  
10 sequences with promoter activity (see, for example, Turgeon et al., *Mol. Cell Biol.*, 1987, 7:3297-3305; Weltring, *Curr. Genet.*, 1995, 28:190-196). This approach provides an easy method for isolating promoter fragments from a wide variety of fungi.

The constructs of the invention also preferably include a terminator  
15 sequence located 3' to the chimeric transcription factor coding sequence. Terminator sequences which function in numerous fungi are known in the art. These include those from *Aspergillus nidulans trpC* (Punt et al., *supra*; Hunter et al., *supra*; Glumoff et al., *supra*), *Lentinus edodes priA* (Yanai et al., *supra*), *Bremia lactucae Ham34* (Judelson et al., *supra*), and *Aspergillus nidulans argB* (Carrez et al., *supra*).

## 20 Construction of Chimeric Transcription Factors

The pre-activated transcription factors of the invention display 1) increased binding, either direct or indirect, to a specific DNA sequence located in a gene regulatory region (e.g., a promoter) *in vivo*, and/or 2) increased transcription activating properties, relative to the precursor molecule. To this  
25 end, it is preferable that part or all of the DBD, the domain of the parental transcription factor which recognizes and binds to the DNA sequences, remain intact. Additional sequences from the parental transcription factor may also remain in the chimeric construct, or they may be removed. The TAD of the



-13-

parental transcription factor may be removed, as the chimeric transcription factor will contain a TAD from another protein, such as the herpesvirus transactivator VP16, as described herein. The TAD from the parental transcription factor may also remain in the chimeric construct.

5           As described above, TADs can be acidic, glutamine-rich, or proline-rich. The ability of each of these TADs to function in any given fungal strain will vary. The acidic TADs have been shown to function in a wide variety of organisms, from *C. elegans* to humans, including fungi. Glutamine-rich and proline-rich TADs have also been shown to function in disparate organisms,  
10 including fungi. As described above, increased transactivation activity may be achieved by using multiple TADs from one category (Tanaka and Herr, *supra*). Furthermore, TADs from more than one class may be used in one chimeric protein (Schwechheimer et al., *supra*; Tanaka et al., *supra*). In the example described below, 4 VP16 TADs and a proline-rich TAD are placed in series.

15           The production of chimeric transcription factors which activate transcription is not limited to the use of parental transcription factors that themselves are transcriptional activators. Using this method, transcription factors which are transcriptional repressors may be converted to transcriptional activators by the addition of a TAD. An example is the *Saccharomyces*  
20 *cerevisiae* Mig1, which is a repressor of *SUC2* expression. Deletion of *mig1* derepresses *SUC2* expression. A chimeric protein in which the DBD of Mig1 is fused to the VP16 TAD can activate transcription from promoters containing Mig1-binding sites and leads to increased expression of *SUC2* (Ostling et al., *Mol Cell Biol.*, 1996, 16:753-61). Thus, the formation of a chimeric  
25 transcriptional activator may be performed for any transcription factor, whether it be an activator or a repressor.

The choice of parental transcription factor for use in the present

-14-

invention depends upon the desired product one wishes to produce. The transcription factor must recognize a sequence in the promoter of a gene of interest. This gene may encode a protein which itself is a desired product, or one which acts as an element (e.g., an enzyme) in the pathway by which a desired product is made by the host fungal cell. For example, a chimeric transcription factor including PacC may be used if the desire is to increase the production of beta-lactam antibiotics. This is achieved by increasing the expression of at least two genes, *ipnA* and *acvA*, which encode enzymes in the penicillin production process.

One skilled in the art will recognize that there are standard techniques, including the ones described herein, which allow for rapid selection and screening of chimeric transcription factor constructs in order to ascertain which transcription factors are the strongest transcriptional activators.

#### Construction of Fungal Expression Vectors

To achieve high expression of the chimeric transcription factor, several types of expression vectors are known in the art (e.g., those described herein). The choice of expression vectors may depend on the type of fungus to be used. For example, expression of a chimeric transcription factor in *Aspergillus nidulans* may be achieved using the *amdS* promoter system (Turnbull et al., *supra*). The promoter element may be modified such that it also contains a DNA sequence recognized by the chimeric transcription factor. The expression of the chimeric transcription factor will induce increased activation from its own promoter, thus amplifying its own production. The expression vector may also include terminator sequences, as described above. For example, a suitable terminator for *Aspergillus nidulans* is the *argB* terminator.

-15-

The vector, once transformed into a fungal cell as described herein, may remain episomal, in which case the vector may also have an origin of replication. The vector may also integrate into the chromosomal DNA of the host cell. The expression of the integrated expression construct may depend on positional effects, and, thus, it may be necessary to screen through or select for transformants to isolate those with suitably high expression. Methods for screening and selection are described herein. The integrated expression construct may also alter the expression of endogenous genes of the fungal cell. This altered expression may be beneficial or detrimental to the survival of the cell or to the purpose of the production of the fungal cell. For example, if the purpose is to increase production of a beta-lactam antibiotic, then loss of expression of *ipnA* (which encodes isopenicillin N-synthase and is required for beta-lactam production) following integration of the expression construct would negate any benefits resulting from expression of the chimeric transcription factor. Thus, a secondary screen of transformants displaying characteristics suitably for the designed purpose may be performed. Methods for determining metabolite production are described herein.

In some cases, it may be beneficial to use a transcription factor which is not chimeric. Overexpression of a parental transcription factor may lead to an increase in secondary metabolites. This overexpressed protein may be constitutively active, due to overexpression or genetic mutation, or it may be regulated in a manner similar to the endogenous transcription factor. The fungal cell may be a wild-type strain, or it may contain one or more mutations (which may also increase production of secondary metabolites). Example mutations include those which result in duplication or rearrangement of biosynthetic genes (e.g., the penicillin gene cluster of *ipnA*, *acvA*, and *aatA*). Reporter genes, such as those described herein, or other exogenous genes may

-16-

also be present in the fungal cells, either episomally or chromosomally.

### Transformation

- In order to introduce the construct into a fungal cell, one may utilize any of numerous transformation protocols (for review, see Punt and van den
- 5 Hondel, *Methods Enzymol.*, 1992, 216:447-457; Timberlake and Marshall, *Science*, 1989, 244:1313-1317; Fincham, *Microbiol. Rev.*, 1989, 53:148-170). Suitable DNA transformation techniques include electroporation, polyethylene glycol-mediated, lithium acetate-mediated, and biolistic transformation (Brown et al., *Mol. Gen. Genet.*, 1998, 259:327-335; Zapanta et al., *Appl. Environ.*
- 10 *Microbiol.*, 1998; 64:2624-2629; Thompson et al., *Yeast*, 1998, 14:565-571; Barreto et al., *FEMS Microbiol. Lett.*, 1997, 156:95-99; Nicolaisen and Geisen, *Microbiol. Res.*, 1996, 151:281-284; Wada et al., *Appl. Microbiol. Biotechnol.*, 1996, 45:652-657; Ozeki et al., *Biosci. Biotechnol. Biochem.*, 1994, 58:2224-2227; Lorito et al., *Curr. Genet.*, 1993, 24:349-356; Oda and Tonomura, *Curr.*
- 15 *Genet.*, 1995, 27:131-134). If desired, one may target the DNA construct to a particular locus. Targeting homologous recombination techniques are currently practiced in many fungi, including, but not limited to, *Candida albicans* (Fonzi and Irwin, *Genetics*, 1993, 134: 717-728), *Ustilago maydis* (Fotheringham and Hollman, *Mol. Cell Biol.*, 1989, 9:4052-4055; Bolker et al., *Mol. Gen. Genet.*,
- 20 1995, 248:547-552), *Yarrowia lipolytica* (Neuveglise et al., *Gene* 1998, 213:37-46; Chen et al., *Appl. Microbiol. Biotechnol.*, 1997, 48:232-235; Cordero et al., *Appl. Microbiol. Biotechnol.*, 1996, 46:143-148), *Acremonium chrysogenum* (Skatrud et al., *Curr. Genet.*, 1987, 12:337-348; Walz and Kuck, *Curr. Genet.*, 1993, 24:421-427), *Magnaporthe grisea* (Sweigard et al., *Mol. Gen. Genet.*,
- 25 1992, 232:183-190); Kershaw et al., *EMBO J.*, 1998, 17:3838-3849), *Histoplasma capsulatum* (Woods et al., *J. Bacteriol.*, 1998, 180:5135-5143)

-17-

and *Aspergillus* sp. (Miller et al., *Mol. Cell Biol.*, 1985, 5:1714-1721; de Ruiter-Jacobs et al., *Curr. Genet.*, 1989, 16:159-163; Gouka et al., *Curr. Genet.*, 1995, 27:536-540; van den Hombergh et al., *Mol. Gen. Genet.*, 1996, 251:542-550; D'Enfert, *Curr. Genet.*, 1996, 30:76-82; Weidner et al., *Curr. Genet.*, 1998, 33:378-385).

### Methods for Selection and Screening Transformants

Reporter genes are useful for isolating transformants expressing functional chimeric transcription factors. The reporter genes may be operably linked to promoter sequence which is regulated by the chimeric transcription factor. Reporter genes include, but are not limited to, genes encoding  $\beta$ -galactosidase (*lacZ*),  $\beta$ -glucoronidase (*GUS*),  $\beta$ -glucosidase, and invertase, amino acid biosynthetic genes, e.g., the yeast *LEU2*, *HIS3*, *LYS2*, *TRP1* genes (or homologous genes from other fungi, such as filamentous fungi, that encode proteins with the similar functional activities), nucleic acid biosynthetic genes, e.g., the yeast *URA3* and *ADE2* genes (or homologous genes from other fungi, such as filamentous fungi, that encode proteins with the similar functional activities), the mammalian chloramphenicol transacetylase (CAT) gene, or any surface antigen gene for which specific antibodies are available. A reporter gene may encode a protein detectable by luminescence or fluorescence, such as green fluorescent protein (GFP). Reporter genes may encode also any protein that provides a phenotypic marker, for example, a protein that is necessary for cell growth or viability, or a toxic protein leading to cell death, or the reporter gene may encode a protein detectable by a color assay leading to the presence or absence of color.

The choice of reporter gene will depend on the type of fungal cell to be transformed. It is preferable to have two reporter genes within the fungal

-18-

cell. One reporter gene, when expressed, may provide a growth advantage to transformed cells which are expressing the chimeric transcription factor. This allows for isolation of such transformants through selective pressures. The other reporter gene may provide a colorimetric marker, such as the *lacZ* gene and its encoded protein,  $\beta$ -galactosidase. Alternatively, the second reporter may provide a fluorescent or luminescent marker, such as GFP. These reporters provide a method of quantifying expression levels from expression constructs comprising chimeric transcription factors. Screens and selections similar to the ones described may be used to optimize construction of chimeric transcription factors or expression constructs.

### Example

The following example describes a method for increasing the level of PacC activity over that caused by proteolysis or specific truncations. This invention may facilitate the increased production of fungal secondary metabolites including, but not limited to, penicillins and cephalosporins. Similar genetic engineering can be performed to alter the function of other transcription factors.

A construct that encodes a chimeric transcription factor is described below. In this example, a proline-rich TAD followed by multiple copies of the acidic-rich TAD from the herpes simplex virus VP16 protein are fused to a truncated, pre-activated PacC from *Aspergillus nidulans* (SEQ ID NO: 7). This construct may be integrated at the *pyrG* locus in *Aspergillus nidulans*, as described below. Expression of this chimeric polypeptide is regulated by the strong PGK promoter from *Aspergillus nidulans* and terminator sequences from the *crnA* gene of *Aspergillus nidulans*.

-19-

Several DNA cloning steps are required to create this chimeric construct. Bluescript KS (Stratagene, LaJolla, CA) is used as a cloning vector. The primers 5'- aactgcagTAGTTGACCGTGTGATTGGGTTCT -3' (SEQ ID NO: 8) (lowercase letters denote sequences introduced for cloning and restriction sites are underlined) and 5'-

5 ccggaattcTTTGTAAGTGGCTTGAAGAT -3' (SEQ ID NO: 9) are used to amplify 347bp of *crnA* terminator sequence from genomic DNA template. The PCR product is *Pst*I/*Eco*RI digested and then cloned into the KS polylinker to produce p1. Subsequently, complementary oligonucleotides 5' -

10 gatccCCCCCCCCCTCCTCCACCCCCACCCCCTCCC -3' (SEQ ID NO: 10) and 5'- GGGAGGGGGTGGGGGTGGAGGAGGGGGGGGg-3' (SEQ ID NO: 11) are annealed (this double-stranded oligonucleotide encodes a proline-rich motif) and the double-stranded product is ligated into *Sma*I/*Bam*HI digested p1, yielding p2.

15 Next, the oligonucleotide primers 5'-

cgcgatcAAAGTCGCCCCCGACCGAT -3' (SEQ ID NO: 12) and 5'-

cgcgatcCCCACCGTACTCGTCAATTCC -3' (SEQ ID NO: 13) are used in PCR reactions to amplify a 258bp fragment using pVP16 (Clontech, Palo Alto, CA) as template. This product encodes the acidic-rich domain of VP16. The

20 product is digested with *Eco*RV, and ligation reaction is performed with >20 fold excess of *Eco*RV insert relative to *Sma*I-digested calf-alkaline phosphatase treated p2. Bacterial transformants are screened for plasmids that contain multiple tandem insertions of VP16 sequence. *Sma*I sites within the VP16 coding sequence allow for determination of the orientation of the insertion.

25 Plasmids are selected that contain four insertions of the VP16 acidic-rich domain (p3). p3, then, encodes a proline-rich domain in-frame with four

-20-

reiterations of the VP16 domain, and these TADs are linked to the *crnA* terminator.

In the next cloning step a truncated form of *pacC* is fused to the coding sequence for the TADs. Primers 5'-

- 5' tgctctagaGGCGCCATGGCCGAAGAAGCG -3' (SEQ ID NO: 14) and 5'-  
cgcggatccGTAACCAGAAGTCATACCGTC -3' (SEQ ID NO: 15) are used to  
amplify a 1419bp product (SEQ ID NO: 16) from an *Aspergillus nidulans*  
cDNA library. This product is *XbaI/BamHI* digested and ligated into digested  
p3 to produce p4. This cloning reaction introduces a form of *pacC* that lacks  
10 the carboxy-terminal 209 amino acids in-frame with the described TADs.

- An additional cloning step is required in order to place the coding  
sequence for this chimera under the control of a strong promoter. Primers 5'-  
ataagaatgcggccgcCCTCTGCATTATTGTCTTATC -3' (SEQ ID NO: 17) and  
5'- tgctctagaAGACATTGTTGCTATAGCTGT -3' (SEQ ID NO: 18) are used  
15 to amplify 689bp of PGK promoter sequence (SEQ ID NO: 19) from  
*Aspergillus nidulans* genomic DNA. This fragment is *NotI/XbaI* digested and  
cloned into digested p4 in order to yield p5. Thus, p5 contains coding sequence  
for an 815 amino acid chimeric transcription factor to be expressed from the  
PGK promoter.

- 20 To decrease the extent of position effects, the p5 construct is targeted  
to the *pyrG* locus. Oligonucleotides 5'-  
tccccgcggATGGAAGCTTCGTTAAGGATAATT-3' (SEQ ID NO: 20) and 5'-  
ataagaatgcggccgcCTACCAGATTAGGGAGCATAT-3' (SEQ ID NO: 21) are  
used to amplify a 2240bp product (SEQ ID NO: 22) from *Aspergillus nidulans*  
25 genomic DNA; this product contains coding and regulatory sequence for the  
*pyrG* gene that encodes orotidine-5'- phosphate decarboxylase. The 2240bp  
fragment is *SacII/NotI* digested, and then cloned into p5 to produce p6; this



-21-

fragment is also cloned into KS to yield p7 (a control construct, containing regulatory sequence for the *pyrG* gene, but no PGK promoter or transcription factor). p6 and p7 are vector that can complement uridine auxotrophy, allowing for selection, and target the chimeric transcription factor to the *pyrG* locus. In addition, primers 5'- tgctctagaGGCGCCATGGCCGAAGAAGCG -3' (SEQ ID NO: 23) and 5' tcccccgggGTAACCAGAAGTCATACCGTC -3' (SEQ ID NO: 24) are used to amplify the truncated form of PacC from an *Aspergillus nidulans* cDNA library. This fragment can be cloned into *XbaI/SmaI* digested p6 to produce p8. p8 is a control construct, used to monitor the activity of pre-activated PacC expressed from the PGK promoter, independent of the presence of heterologous TADs.

PEG- $\text{CaCl}_2$  (or other methods, described herein) may be used to transform protoplasts of a uridine auxotroph carrying a *pyrG* mutation (Ballance and Turner, *Gene*, 1985, 36:321-331). p6, p7, and p8 plasmid DNA are used to transform to uridine prototrophy. PCR and Southern analysis are performed to confirm single-copy integration at *pyrG*.

Several methods may be employed to assess the activity of wild-type, pre-activated, and chimeric PacC-TAD factors. Samples of mycelia may be taken from parallel fermentation of strains containing p6, p7, and p8. Northern blot analysis may be performed on RNA prepared from extracts of these mycelia. Probes are prepared from coding sequence for the *ipnA* and *acvA* genes of *Aspergillus nidulans*. Reporter constructs are valuable tools for examining the level of PacC activation. For example, *ipnA* and *acvA* are divergently transcribed from a common regulatory sequence. One may use constructs (e.g., pAXB4A; Brakhage et al., *supra*) that contain *ipnA-lacZ* and *acvA-uidA* reporters within the same plasmid; this particular plasmid can be targeted to the *argB* locus to ensure integration at a specific genomic locus. A

-22-

strain carrying both *argB* and *pyrG* mutations can be sequentially transformed with the *pyrG* and reporter vectors, and enzyme assays can be performed on extracts from mycelia (van Gorcom et al., *Gene*, 1985, 40:99-106; Pobjecky et al., *Mol. Gen. Genet.*, 1990, 220:314-316). In addition, bioassays can be done  
5 to determine whether chimeric transcription factors increase the production of fungal secondary metabolites such as penicillin. Supernatant fluid from fermentations can be centrifuged and applied to wells containing indicator organisms such as *Bacillus calidolactis* (Smith et al., *Mol. Gen. Genet.*, 1989, 216:492-497). The application of all of these methods will promote a rapid and  
10 quantitative analysis of the efficacy of chimeric transcription factors.

#### Enhancement of Secondary Metabolite Production

The constructs and methods described herein may be used to increase the yields of currently marketed pharmaceuticals whose production, in whole or in part, is dependent upon a fungal fermentation. For example, in *Aspergillus*  
15 *nidulans*, penicillin biosynthesis is catalyzed by three enzymes encoded by *ipnA*, *acvA*, and *aatA*. Two of these genes, *ipnA* and *acvA*, are regulated directly by PacC. For example, *P<sub>ipnA</sub>* contains at least three PacC binding sites (*ipnA2*, *ipnA3*, and *ipnA4AB*)(Espeso and Penalva, *J. Biol. Chem.*, 1996, 271:28825-28830). Expression of a truncated form of PacC has been shown to  
20 increase both expression of *ipnA* and *acvA* as well as production of penicillin. Activation (i.e., proteolytic cleavage) of PacC requires the proteins encoded by the *palA*, *palB*, *palC*, *palF*, *palH*, and *palI* genes. It is possible that increased expression of at least some of these genes would result in increased production of penicillin. In the example described herein, *ipnA* and *acvA* expression are  
25 targeted for increase by formation of a chimeric transcription factor including the DNA-binding domain of PacC and 4 VP16 acidic TADs and a proline-rich

-23-

TAD. Using the methods of the invention, production of other secondary metabolites can also be increased.

Examples of marketed secondary metabolites whose yields during fermentation could be increased by the methods of the invention include, without limitation, cyclosporin, penicillin, cephalosporin, ergot alkaloids, lovastatin, mevastatin, and the biosynthetic intermediates thereof. In addition, such methods can also be used to increase the likelihood of identifying new secondary metabolites with medicinal or agricultural value by increasing the concentration of such metabolites (and hence, the likelihood of detection by chemical or bioassay) in a fermentation broth.

#### Production and Detection Methods for Fungal Secondary Metabolites

Methods for fermentation and production of beta-lactam antibiotics, statins, ergot alkaloids, cyclosporin, and other fungal metabolites are described in Masurekar (*Biotechnology*, 1992, 21: 241-301), and references therein. The detection of secondary metabolites is specific for each metabolite and well-known to those practiced in the art. General methods to assess production and integrity of compounds in fermentation broths include, but are not limited to, bioassays for antimicrobial activity, high-performance liquid chromatography (HPLC) analysis, nuclear magnetic resonance, thin-layer chromatography, and absorbance spectrometry. Purification of metabolites from a fermentation broth can include removal of fungal cells or hyphae by centrifugation or filtration, adjustment of pH and/or salt concentrations after fermentation (to enhance solubility and/or subsequent extraction efficiency), and extraction of broths with appropriate organic solvents.

What is claimed is:

-24-

1. A chimeric transcription factor comprising
  - (a) a pre-activated transcription factor functional in a fungal strain,  
and
  - (b) a transcription activation domain that is different from the  
5 transcription activation domain naturally associated with said transcription  
factor.
2. The chimeric transcription factor of claim 1, wherein said  
chimeric transcription factor activates transcription in a manner greater than  
said pre-activated transcription factor.
- 10 3. The chimeric transcription factor of claim 1, wherein said pre-  
activated transcription factor is pre-activated by truncation.
4. The chimeric transcription factor of claim 1, wherein said pre-  
activated transcription factor comprises a substitution of a serine or threonine  
residue with an alanine, aspartic acid, or glutamic acid residue, wherein said  
15 substitution pre-activates said transcription factor.
5. The chimeric transcription factor of claim 3, wherein said pre-  
activated transcription factor is substantially identical to *Aspergillus nidulans*  
PacC.
6. A chimeric transcription factor comprising
  - (a) a transcription factor substantially identical to *Aspergillus*  
20 *nidulans* PacC, and

-25-

(b) a transcription activation domain that is different from the transcription activation domain naturally associated with said transcription factor.

7. The chimeric transcription factor of claim 1, wherein said pre-activated transcription factor comprises amino acid sequence shown in Fig. 1 (SEQ ID Nos: 1-6).

8. The chimeric transcription factor of claim 1, wherein said pre-activated transcription factor binds to a DNA sequence comprising 5'-GCCAAG-3' or 5'-GCCAGG-3'.

9. A vector comprising DNA encoding the chimeric transcription factor of claim 1 operably linked to a promoter capable of controlling expression of said chimeric transcription factor in a fungal strain.

10. A fungal cell that contains and expresses the DNA of claim 9.

11. The fungal cell of claim 10, wherein said fungal cell is a filamentous fungal cell.

12. The fungal cell of claim 10, wherein said cell produces a secondary metabolite and wherein expression of said DNA increases the production of said secondary metabolite by said cell.

13. The fungal cell of claim 12, wherein said secondary metabolite is non-proteinaceous.

-26-

14. The fungal cell of claim 12, wherein said secondary metabolite is a protein or peptide.

15. A method of producing a secondary metabolite, said method comprising culturing the fungal cell of claim 10 under secondary metabolite-producing conditions.

16. A method of producing a secondary metabolite, said method comprising the steps of

(a) introducing into a fungal cell a vector comprising (i) a promoter capable of controlling gene expression in said fungal cell, and (ii) a nucleic acid encoding a transcription factor comprising (i) a DNA-binding domain and (ii) a transcription activation domain; and

(b) culturing said fungal cell under secondary metabolite-producing conditions.

17. The method of claim 16, wherein said fungal cell is a filamentous fungal cell.

18. The method of claim 16, wherein said transcription factor is a chimeric transcription factor.

19. The method of claim 16, wherein said transcription factor is a pre-activated transcription factor.

20. The method of claim 16, wherein said transcription factor is pre-activated by substitution of a serine or threonine residue with an alanine,

-27-

aspartic acid, or glutamic acid residue, wherein said substitution pre-activates said transcription factor.

21. The method of claim 16, wherein said transcription factor is pre-activated by truncation.

5           22. The method of claim 16, wherein said DNA binding domain is from a fungal transcriptional activator.

23. The method of claim 16, wherein said DNA binding domain is from a fungal transcriptional repressor.

10           24. The method of claim 16, wherein said transcription activation domain that is different from the transcription activation domain naturally associated with said transcription factor.

Figure 1  
Sheet 1 of 2



2/2

A. nidulans	311	ONLSLPVLTAAAPLPEYQAMPAVAVAA..SGPYG..GGPHPAAYHLPPMSNVRTKN..DLIN
A. niger	312	QSLSLPVLSSGAPLPEYQPMPPAVAVG..GGGYSPGGAPSAAYHLPPMSNVRTKN..DLIN
P. chrysogenum	313	QALQLPFLS-GPAPPEYQOMPAVAVGGGGGYG..GGGAPGPAYHLPPMSNVRTKN..DLIN
Y. lipolytica	314	EQYIGLIGGQ..-GQAQASPTPOTAT..TTSATPAPAAAPHOATPPQQQLPSFKQGDYQE
C. albicans	315	KRMKAAGTEY..-NLDVFNKLN..HLDHLLHHHPQQOH..QQQYGGN..LYYE
S. cerevisiae	316	NSRSMNVYE DQ..-CSNKTIANATQ FTKLSRNMNTNNVITLQQSGGSTESSISSGRIPVA
A. nidulans	317	..
A. niger	318	..
P. chrysogenum	319	..
Y. lipolytica	320	..
C. albicans	321	..
S. cerevisiae	322	..
A. nidulans	323	..
A. niger	324	..
P. chrysogenum	325	..
Y. lipolytica	326	..
C. albicans	327	..
S. cerevisiae	328	..
A. nidulans	329	..
A. niger	330	..
P. chrysogenum	331	..
Y. lipolytica	332	..
C. albicans	333	..
S. cerevisiae	334	..
A. nidulans	335	..
A. niger	336	..
P. chrysogenum	337	..
Y. lipolytica	338	..
C. albicans	339	..
S. cerevisiae	340	..
A. nidulans	341	..
A. niger	342	..
P. chrysogenum	343	..
Y. lipolytica	344	..
C. albicans	345	..
S. cerevisiae	346	..
A. nidulans	347	..
A. niger	348	..
P. chrysogenum	349	..
Y. lipolytica	350	..
C. albicans	351	..
S. cerevisiae	352	..
A. nidulans	353	..
A. niger	354	..
P. chrysogenum	355	..
Y. lipolytica	356	..
C. albicans	357	..
S. cerevisiae	358	..
A. nidulans	359	..
A. niger	360	..
P. chrysogenum	361	..
Y. lipolytica	362	..
C. albicans	363	..
S. cerevisiae	364	..
A. nidulans	365	..
A. niger	366	..
P. chrysogenum	367	..
Y. lipolytica	368	..
C. albicans	369	..
S. cerevisiae	370	..
A. nidulans	371	..
A. niger	372	..
P. chrysogenum	373	..
Y. lipolytica	374	..
C. albicans	375	..
S. cerevisiae	376	..
A. nidulans	377	..
A. niger	378	..
P. chrysogenum	379	..
Y. lipolytica	380	..
C. albicans	381	..
S. cerevisiae	382	..
A. nidulans	383	..
A. niger	384	..
P. chrysogenum	385	..
Y. lipolytica	386	..
C. albicans	387	..
S. cerevisiae	388	..
A. nidulans	389	..
A. niger	390	..
P. chrysogenum	391	..
Y. lipolytica	392	..
C. albicans	393	..
S. cerevisiae	394	..
A. nidulans	395	..
A. niger	396	..
P. chrysogenum	397	..
Y. lipolytica	398	..
C. albicans	399	..
S. cerevisiae	400	..
A. nidulans	401	..
A. niger	402	..
P. chrysogenum	403	..
Y. lipolytica	404	..
C. albicans	405	..
S. cerevisiae	406	..
A. nidulans	407	..
A. niger	408	..
P. chrysogenum	409	..
Y. lipolytica	410	..
C. albicans	411	..
S. cerevisiae	412	..
A. nidulans	413	..
A. niger	414	..
P. chrysogenum	415	..
Y. lipolytica	416	..
C. albicans	417	..
S. cerevisiae	418	..
A. nidulans	419	..
A. niger	420	..
P. chrysogenum	421	..
Y. lipolytica	422	..
C. albicans	423	..
S. cerevisiae	424	..
A. nidulans	425	..
A. niger	426	..
P. chrysogenum	427	..
Y. lipolytica	428	..
C. albicans	429	..
S. cerevisiae	430	..
A. nidulans	431	..
A. niger	432	..
P. chrysogenum	433	..
Y. lipolytica	434	..
C. albicans	435	..
S. cerevisiae	436	..
A. nidulans	437	..
A. niger	438	..
P. chrysogenum	439	..
Y. lipolytica	440	..
C. albicans	441	..
S. cerevisiae	442	..
A. nidulans	443	..
A. niger	444	..
P. chrysogenum	445	..
Y. lipolytica	446	..
C. albicans	447	..
S. cerevisiae	448	..
A. nidulans	449	..
A. niger	450	..
P. chrysogenum	451	..
Y. lipolytica	452	..
C. albicans	453	..
S. cerevisiae	454	..
A. nidulans	455	..
A. niger	456	..
P. chrysogenum	457	..
Y. lipolytica	458	..
C. albicans	459	..
S. cerevisiae	460	..
A. nidulans	461	..
A. niger	462	..
P. chrysogenum	463	..
Y. lipolytica	464	..
C. albicans	465	..
S. cerevisiae	466	..
A. nidulans	467	..
A. niger	468	..
P. chrysogenum	469	..
Y. lipolytica	470	..
C. albicans	471	..
S. cerevisiae	472	..
A. nidulans	473	..
A. niger	474	..
P. chrysogenum	475	..
Y. lipolytica	476	..
C. albicans	477	..
S. cerevisiae	478	..
A. nidulans	479	..
A. niger	480	..
P. chrysogenum	481	..
Y. lipolytica	482	..
C. albicans	483	..
S. cerevisiae	484	..
A. nidulans	485	..
A. niger	486	..
P. chrysogenum	487	..
Y. lipolytica	488	..
C. albicans	489	..
S. cerevisiae	490	..
A. nidulans	491	..
A. niger	492	..
P. chrysogenum	493	..
Y. lipolytica	494	..
C. albicans	495	..
S. cerevisiae	496	..
A. nidulans	497	..
A. niger	498	..
P. chrysogenum	499	..
Y. lipolytica	500	..
C. albicans	501	..
S. cerevisiae	502	..
A. nidulans	503	..
A. niger	504	..
P. chrysogenum	505	..
Y. lipolytica	506	..
C. albicans	507	..
S. cerevisiae	508	..
A. nidulans	509	..
A. niger	510	..
P. chrysogenum	511	..
Y. lipolytica	512	..
C. albicans	513	..
S. cerevisiae	514	..
A. nidulans	515	..
A. niger	516	..
P. chrysogenum	517	..
Y. lipolytica	518	..
C. albicans	519	..
S. cerevisiae	520	..
A. nidulans	521	..
A. niger	522	..
P. chrysogenum	523	..
Y. lipolytica	524	..
C. albicans	525	..
S. cerevisiae	526	..
A. nidulans	527	..
A. niger	528	..
P. chrysogenum	529	..
Y. lipolytica	530	..
C. albicans	531	..
S. cerevisiae	532	..
A. nidulans	533	..
A. niger	534	..
P. chrysogenum	535	..
Y. lipolytica	536	..
C. albicans	537	..
S. cerevisiae	538	..
A. nidulans	539	..
A. niger	540	..
P. chrysogenum	541	..
Y. lipolytica	542	..
C. albicans	543	..
S. cerevisiae	544	..
A. nidulans	545	..
A. niger	546	..
P. chrysogenum	547	..
Y. lipolytica	548	..
C. albicans	549	..
S. cerevisiae	550	..
A. nidulans	551	..
A. niger	552	..
P. chrysogenum	553	..
Y. lipolytica	554	..
C. albicans	555	..
S. cerevisiae	556	..
A. nidulans	557	..
A. niger	558	..
P. chrysogenum	559	..
Y. lipolytica	560	..
C. albicans	561	..
S. cerevisiae	562	..
A. nidulans	563	..
A. niger	564	..
P. chrysogenum	565	..
Y. lipolytica	566	..
C. albicans	567	..
S. cerevisiae	568	..
A. nidulans	569	..
A. niger	570	..
P. chrysogenum	571	..
Y. lipolytica	572	..
C. albicans	573	..
S. cerevisiae	574	..
A. nidulans	575	..
A. niger	576	..
P. chrysogenum	577	..
Y. lipolytica	578	..
C. albicans	579	..
S. cerevisiae	580	..
A. nidulans	581	..
A. niger	582	..
P. chrysogenum	583	..
Y. lipolytica	584	..
C. albicans	585	..
S. cerevisiae	586	..
A. nidulans	587	..
A. niger	588	..
P. chrysogenum	589	..
Y. lipolytica	590	..
C. albicans	591	..
S. cerevisiae	592	..
A. nidulans	593	..
A. niger	594	..
P. chrysogenum	595	..
Y. lipolytica	596	..
C. albicans	597	..
S. cerevisiae	598	..
A. nidulans	599	..
A. niger	600	..
P. chrysogenum	601	..
Y. lipolytica	602	..
C. albicans	603	..
S. cerevisiae	604	..
A. nidulans	605	..
A. niger	606	..
P. chrysogenum	607	..
Y. lipolytica	608	..
C. albicans	609	..
S. cerevisiae	610	..
A. nidulans	611	..
A. niger	612	..
P. chrysogenum	613	..
Y. lipolytica	614	..
C. albicans	615	..
S. cerevisiae	616	..
A. nidulans	617	..
A. niger	618	..
P. chrysogenum	619	..
Y. lipolytica	620	..
C. albicans	621	..
S. cerevisiae	622	..
A. nidulans	623	..
A. niger	624	..
P. chrysogenum	625	..
Y. lipolytica	626	..
C. albicans	627	..
S. cerevisiae	628	..
A. nidulans	629	..
A. niger	630	..
P. chrysogenum	631	..
Y. lipolytica	632	..
C. albicans	633	..
S. cerevisiae	634	..
A. nidulans	635	..
A. niger	636	..
P. chrysogenum	637	..
Y. lipolytica	638	..
C. albicans	639	..
S. cerevisiae	640	..
A. nidulans	641	..
A. niger	642	..
P. chrysogenum	643	..
Y. lipolytica	644	..
C. albicans	645	..
S. cerevisiae	646	..
A. nidulans	647	..
A. niger	648	..
P. chrysogenum	649	..
Y. lipolytica	650	..
C. albicans	651	..
S. cerevisiae	652	..
A. nidulans	653	..
A. niger	654	..
P. chrysogenum	655	..
Y. lipolytica	656	..
C. albicans	657	..
S. cerevisiae	658	..
A. nidulans	659	..
A. niger	660	..
P. chrysogenum	661	..
Y. lipolytica	662	..
C. albicans	663	..
S. cerevisiae	664	..
A. nidulans	665	..
A. niger	666	..
P. chrysogenum	667	..
Y. lipolytica	668	..
C. albicans	669	..
S. cerevisiae	670	..
A. nidulans	671	..
A. niger	672	..
P. chrysogenum	673	..
Y. lipolytica	674	..
C. albicans	675	..
S. cerevisiae	676	..
A. nidulans	677	..
A. niger	678	..
P. chrysogenum	679	..
Y. lipolytica	680	..
C. albicans	681	..
S. cerevisiae	682	..
A. nidulans	683	..
A. niger	684	..
P. chrysogenum	685	..
Y. lipolytica	686	..
C. albicans	687	..
S. cerevisiae	688	..
A. nidulans	689	..
A. niger	690	..
P. chrysogenum	691	..
Y. lipolytica	692	..
C. albicans	693	..
S. cerevisiae	694	..
A. nidulans	695	..
A. niger	696	..
P. chrysogenum	697	..
Y. lipolytica	698	..
C. albicans	699	..
S. cerevisiae	700	..
A. nidulans	701	..
A. niger	702	..
P. chrysogenum	703	..
Y. lipolytica	704	..
C. albicans	705	..
S. cerevisiae	706	..
A. nidulans	707	..
A. niger	708	..
P. chrysogenum	709	..
Y. lipolytica	710	..
C. albicans	711	..
S. cerevisiae	712	..
A. nidulans	713	..
A. niger	714	..
P. chrysogenum	715	..
Y. lipolytica	716	..
C. albicans	717	..
S. cerevisiae	718	..
A. nidulans	719	..
A. niger	720	..
P. chrysogenum	721	..
Y. lipolytica	722	..
C. albicans	723	..
S. cerevisiae	724	..
A. nidulans	725	..
A. n		

## SEQUENCE LISTING

&lt;110&gt; Microbia, Inc.

<120> CHIMERIC PRE-ACTIVATED TRANSCRIPTION  
FACTORS

&lt;130&gt; 50078/004WO2

&lt;150&gt; 60/066,129

&lt;151&gt; 1997-11-19

&lt;150&gt; 60/066,308

&lt;151&gt; 1997-11-21

&lt;150&gt; 60/066,462

&lt;151&gt; 1997-11-24

&lt;160&gt; 24

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 678

&lt;212&gt; PRT

&lt;213&gt; Aspergillus nidulans

&lt;400&gt; 1

Met	Leu	Gly	Ala	Met	Ala	Glu	Glu	Ala	Val	Ala	Pro	Val	Ala	Val	Pro
1				5					10					15	
Thr	Thr	Gln	Glu	Gln	Pro	Thr	Ser	Gln	Pro	Ala	Ala	Ala	Gln	Val	Thr
			20					25					30		
Thr	Val	Thr	Ser	Pro	Ser	Val	Thr	Ala	Thr	Ala	Ala	Ala	Ala	Thr	Ala
		35					40					45			
Ala	Val	Ala	Ser	Pro	Gln	Ala	Asn	Gly	Asn	Ala	Ala	Ser	Pro	Val	Ala
	50					55				60					
Pro	Ala	Ser	Ser	Thr	Ser	Arg	Pro	Ala	Glu	Glu	Leu	Thr	Cys	Met	Trp
65					70				75					80	
Gln	Gly	Cys	Ser	Glu	Lys	Leu	Pro	Thr	Pro	Glu	Ser	Leu	Tyr	Glu	His
			85					90					95		
Val	Cys	Glu	Arg	His	Val	Gly	Arg	Lys	Ser	Thr	Asn	Asn	Leu	Asn	Leu
			100				105						110		
Thr	Cys	Gln	Trp	Gly	Ser	Cys	Arg	Thr	Thr	Thr	Val	Lys	Arg	Asp	His
		115				120					125				
Ile	Thr	Ser	His	Ile	Arg	Val	His	Val	Pro	Leu	Lys	Pro	His	Lys	Cys
	130					135				140					
Asp	Phe	Cys	Gly	Lys	Ala	Phe	Lys	Arg	Pro	Gln	Asp	Leu	Lys	Lys	His
145					150					155					160
Val	Lys	Thr	His	Ala	Asp	Asp	Ser	Val	Leu	Val	Arg	Ser	Pro	Glu	Pro
			165						170					175	
Gly	Ser	Arg	Asn	Pro	Asp	Met	Met	Phe	Gly	Gly	Asn	Gly	Lys	Gly	Tyr
			180					185					190		
Ala	Ala	Ala	His	Tyr	Phe	Glu	Pro	Ala	Leu	Asn	Pro	Val	Pro	Ser	Gln
			195				200						205		

Gly Tyr Ala His Gly Pro Pro Gln Tyr Tyr Gln Ala His His Ala Pro  
 210 215 220  
 Gln Pro Ser Asn Pro Ser Tyr Gly Asn Val Tyr Tyr Ala Leu Asn Thr  
 225 230 235 240  
 Gly Pro Glu Pro His Gln Ala Ser Tyr Glu Ser Lys Lys Arg Gly Tyr  
 245 250 255  
 Asp Ala Leu Asn Glu Phe Phe Gly Asp Leu Lys Arg Arg Gln Phe Asp  
 260 265 270  
 Pro Asn Ser Tyr Ala Ala Val Gly Gln Arg Leu Leu Ser Leu Gln Asn  
 275 280 285  
 Leu Ser Leu Pro Val Leu Thr Ala Ala Pro Leu Pro Glu Tyr Gln Ala  
 290 295 300  
 Met Pro Ala Pro Val Ala Val Ala Ser Gly Pro Tyr Gly Gly Gly Pro  
 305 310 315 320  
 His Pro Ala Pro Ala Tyr His Leu Pro Pro Met Ser Asn Val Arg Thr  
 325 330 335  
 Lys Asn Asp Leu Ile Asn Ile Asp Gln Phe Leu Gln Gln Met Gln Asp  
 340 345 350  
 Thr Ile Tyr Glu Asn Asp Asp Asn Val Ala Ala Ala Gly Val Ala Gln  
 355 360 365  
 Pro Gly Ala His Tyr Ile His Asn Gly Ile Ser Tyr Arg Thr Thr His  
 370 375 380  
 Ser Pro Pro Thr Gln Leu Pro Ser Ala His Ala Thr Thr Gln Thr Thr  
 385 390 395 400  
 Ala Gly Pro Ile Ile Ser Asn Thr Ser Ala His Ser Pro Ser Ser Ser  
 405 410 415  
 Thr Pro Ala Leu Thr Pro Pro Ser Ser Ala Gln Ser Tyr Thr Ser Gly  
 420 425 430  
 Arg Ser Pro Ile Ser Leu Pro Ser Ala His Arg Val Ser Pro Pro His  
 435 440 445  
 Glu Ser Gly Ser Ser Met Tyr Pro Arg Leu Pro Ser Ala Thr Asp Gly  
 450 455 460  
 Met Thr Ser Gly Tyr Thr Ala Ala Ser Ser Ala Ala Pro Pro Ser Thr  
 465 470 475 480  
 Leu Gly Gly Ile Phe Asp Asn Asp Glu Arg Arg Arg Tyr Thr Gly Gly  
 485 490 495  
 Thr Leu Gln Arg Ala Arg Pro Ala Ser Arg Ala Ala Ser Glu Ser Met  
 500 505 510  
 Asp Leu Ser Ser Asp Asp Lys Glu Ser Gly Glu Arg Thr Pro Lys Gln  
 515 520 525  
 Ile Ser Ala Ser Leu Ile Asp Pro Ala Leu His Ser Gly Ser Pro Gly  
 530 535 540  
 Glu Asp Asp Val Thr Arg Thr Ala Lys Ala Ala Thr Glu Val Ala Glu  
 545 550 555 560  
 Arg Ser Asp Val Gln Ser Glu Trp Val Glu Lys Val Arg Leu Ile Glu  
 565 570 575  
 Tyr Leu Arg Asn Tyr Ile Ala Asn Arg Leu Glu Arg Gly Glu Phe Ser  
 580 585 590  
 Asp Asp Ser Glu Gln Glu Gln Asp Gln Glu Gln Glu Gln Asp Gln Glu  
 595 600 605  
 Gln Glu Gln Asp Gln Glu Gln Gly Gln Asp Arg Val Ser Arg Ser Pro  
 610 615 620  
 Val Ser Lys Ala Asp Val Asp Met Glu Gly Val Glu Arg Asp Ser Leu  
 625 630 635 640  
 Pro Arg Ser Pro Arg Thr Val Pro Ile Lys Thr Asp Gly Glu Ser Ala

```
<210> 2
<211> 667
<212> PRT
<213> Aspergillus niger
```

3

```

          340                      345                      350
Asn Asp Asp Asn Val Ala Ala Ala Gly Val Ala Gln Pro Gly Ala His
          355                      360                      365
Tyr Val His Gly Gly Met Ser Tyr Arg Thr Thr His Ser Pro Pro Thr
          370                      375                      380
Gln Leu Pro Pro Ser His Ala Thr Ala Thr Ser Ser Ala Ser Met Met
385                      390                      395                      400
Pro Asn Pro Ala Thr His Ser Pro Ser Thr Gly Thr Pro Ala Leu Thr
          405                      410                      415
Pro Pro Ser Ser Ala Gln Ser Tyr Thr Ser Gly Arg Ser Pro Val Ser
          420                      425                      430
Leu Pro Ser Ala Thr Arg Val Ser Pro Pro His His Glu Gly Gly Ser
          435                      440                      445
Met Tyr Pro Arg Leu Pro Ser Ala Thr Met Ala Asp Ser Met Ala Ala
          450                      455                      460
Gly Tyr Pro Thr Ala Ser Ser Thr Ala Pro Pro Ser Thr Leu Gly Gly
465                      470                      475                      480
Ile Phe Asp His Asp Asp Arg Arg Arg Tyr Thr Gly Gly Thr Leu Gln
          485                      490                      495
Arg Ala Arg Pro Glu Thr Arg Gln Leu Ser Glu Glu Met Asp Leu Thr
          500                      505                      510
Gln Asp Ser Lys Asp Glu Gly Glu Arg Thr Pro Lys Ala Lys Glu His
          515                      520                      525
Ser Ser Pro Ser Ser Pro Glu Arg Ile Ser Ala Ser Leu Ile Asp Pro
          530                      535                      540
Ala Leu Ser Gly Thr Ala Ala Glu Ala Glu Ala Thr Leu Arg Thr Ala
545                      550                      555                      560
Gln Ala Ala Thr Glu Val Ala Glu Arg Ala Asp Val Gln Trp Val Glu
          565                      570                      575
Lys Val Arg Leu Ile Glu Tyr Leu Arg Asn Tyr Ile Ala Ser Arg Leu
          580                      585                      590
Glu Arg Gly Glu Phe Glu Asn Asn Glu Ser Gly Gly Gly Asn Ser Ser
          595                      600                      605
Ser Asn Gly Ser Ser His Glu Gln Thr Pro Glu Ala Ser Pro Asp Thr
          610                      615                      620
His Met Glu Gly Val Glu Ser Glu Val Pro Ser Lys Ala Glu Glu Pro
625                      630                      635                      640
Ala Val Lys Pro Glu Ala Gly Asp Val Val Met Tyr Pro Thr Leu Arg
          645                      650                      655
Ala Val Asp Glu Asp Gly Asp Ser Lys Met Pro
          660                      665

```

```

<210> 3
<211> 643
<212> PRT
<213> Penicillium chrysogenum

```

```

<400> 3
Met Thr Glu Asn His Thr Pro Ser Thr Thr Gln Pro Thr Leu Pro Ala
 1                      5                      10                      15
Pro Val Ala Glu Ala Ala Pro Ile Gln Ala Asn Pro Ala Pro Ser Ala
          20                      25                      30
Ser Val Thr Ala Thr Ala Ala Ala Thr Ala Ala Val Asn Asn Ala
          35                      40                      45
Pro Ser Met Asn Gly Ala Gly Glu Gln Leu Pro Cys Gln Trp Val Gly

```

50						55						60					
Cys	Thr	Glu	Lys	Ser	Pro	Thr	Ala	Glu	Ser	Leu	Tyr	Glu	His	Val	Cys		
65					70					75					80		
Glu	Arg	His	Val	Gly	Arg	Lys	Ser	Thr	Asn	Asn	Leu	Asn	Leu	Thr	Cys		
				85					90					95			
Gln	Trp	Gly	Thr	Cys	Asn	Thr	Thr	Thr	Val	Lys	Arg	Asp	His	Ile	Thr		
			100					105					110				
Ser	His	Ile	Arg	Val	His	Val	Pro	Leu	Lys	Pro	His	Lys	Cys	Asp	Phe		
		115					120					125					
Cys	Gly	Lys	Ala	Phe	Lys	Arg	Pro	Gln	Asp	Leu	Lys	Lys	His	Val	Lys		
	130					135				140							
Thr	His	Ala	Asp	Asp	Ser	Glu	Ile	Arg	Ser	Pro	Glu	Pro	Gly	Met	Lys		
145					150					155					160		
His	Pro	Asp	Met	Met	Phe	Pro	Gln	Asn	Pro	Arg	Gly	Ser	Pro	Ala	Ala		
			165						170					175			
Thr	His	Tyr	Phe	Glu	Ser	Pro	Ile	Asn	Gly	Ile	Asn	Gly	Gln	Tyr	Ser		
		180						185					190				
His	Ala	Pro	Pro	Pro	Gln	Tyr	Tyr	Gln	Pro	His	Pro	Pro	Pro	Gln	Ala		
	195					200						205					
Pro	Asn	Pro	His	Ser	Tyr	Gly	Asn	Leu	Tyr	Tyr	Ala	Leu	Ser	Gln	Gly		
	210					215					220						
Gln	Glu	Gly	Gly	His	Pro	Tyr	Asp	Arg	Lys	Arg	Gly	Tyr	Asp	Ala	Leu		
225					230					235				240			
Asn	Glu	Phe	Phe	Gly	Asp	Leu	Lys	Arg	Arg	Gln	Phe	Asp	Pro	Asn	Ser		
			245						250					255			
Tyr	Ala	Ala	Val	Gly	Gln	Arg	Leu	Leu	Gly	Leu	Gln	Ala	Leu	Gln	Leu		
		260					265						270				
Pro	Phe	Leu	Ser	Gly	Pro	Ala	Pro	Glu	Tyr	Gln	Gln	Met	Pro	Ala	Pro		
	275						280					285					
Val	Ala	Val	Gly	Gly	Gly	Gly	Gly	Gly	Tyr	Gly	Gly	Gly	Ala	Pro	Gln		
	290					295					300						
Pro	Pro	Gly	Tyr	His	Leu	Pro	Pro	Met	Ser	Asn	Val	Arg	Thr	Lys	Asn		
305					310					315				320			
Asp	Leu	Ile	Asn	Ile	Asp	Gln	Phe	Leu	Glu	Gln	Met	Gln	Asn	Thr	Ile		
			325						330					335			
Tyr	Glu	Ser	Asp	Glu	Asn	Val	Ala	Ala	Ala	Gly	Val	Ala	Gln	Pro	Gly		
		340					345						350				
Ala	His	Tyr	Val	His	Gly	Gly	Met	Asn	His	Arg	Thr	Thr	His	Ser	Pro		
	355						360					365					
Pro	Thr	His	Ser	Arg	Gln	Ala	Thr	Leu	Leu	Gln	Leu	Pro	Ser	Ala	Pro		
	370					375					380						
Met	Ala	Ala	Ala	Thr	Ala	His	Ser	Pro	Ser	Val	Gly	Thr	Pro	Ala	Leu		
385					390					395				400			
Thr	Pro	Pro	Ser	Ser	Ala	Gln	Ser	Tyr	Thr	Ser	Asn	Arg	Ser	Pro	Ile		
			405						410					415			
Ser	Leu	His	Ser	Ser	Arg	Val	Ser	Pro	Pro	His	Glu	Glu	Ala	Ala	Pro		
			420					425					430				
Gly	Met	Tyr	Pro	Arg	Leu	Pro	Ala	Ala	Ile	Cys	Ala	Asp	Ser	Met	Thr		
	435						440					445					
Ala	Gly	Tyr	Pro	Thr	Ala	Ser	Gly	Ala	Ala	Pro	Pro	Ser	Thr	Leu	Ser		
	450					455					460						
Gly	Ala	Tyr	Asp	His	Asp	Arg	Arg	Arg	Tyr	Thr	Gly	Gly	Thr	Leu			
465					470				475					480			
Gln	Arg	Ala	Arg	Pro	Ala	Glu	Arg	Ala	Ala	Thr	Glu	Asp	Arg	Met	Asp		
			485						490					495			

```

Ile Ser Gln Asp Ser Lys His Asp Gly Glu Arg Thr Pro Lys Ala Met
      500      505      510
His Ile Ser Ala Ser Leu Ile Asp Pro Ala Leu Ser Gly Thr Ser Ser
      515      520      525
Asp Pro Glu Gln Glu Ser Ala Lys Arg Thr Ala Ala Thr Ala Thr Glu
      530      535      540
Val Ala Glu Arg Asp Val Asn Val Ala Trp Val Glu Lys Val Arg Leu
545      550      555      560
Leu Glu Asn Leu Arg Arg Leu Val Ser Gly Leu Leu Glu Ala Gly Ser
      565      570      575
Leu Thr Pro Glu Tyr Gly Val Gln Thr Ser Ser Ala Ser Pro Thr Pro
      580      585      590
Gly Leu Asp Ala Met Glu Gly Val Glu Thr Ala Ser Val Arg Ala Ala
      595      600      605
Ser Glu Gln Ala Arg Glu Glu Pro Lys Ser Glu Ser Glu Gly Val Phe
      610      615      620
Tyr Pro Thr Leu Arg Gly Val Asp Glu Asp Glu Asp Gly Asp Ser Lys
625      630      635      640
Met Pro Glu

```

```

<210> 4
<211> 585
<212> PRT
<213> Yarrowia lipolytica

```

```

<400> 4
Met Ala Ser Tyr Pro Tyr Leu Ala Gln Ser Gln Pro Pro Gln Gln Gln
 1      5      10      15
Gln Gln Gln Gln Gln Gln Pro Gln Gln Gln Ser Gln Gln Leu Pro Thr
      20      25      30
Thr Ala Pro Ser Ala Ala Pro Gln Val Asn Asn Thr Thr Ala Asn Lys
      35      40      45
Pro Leu Tyr Pro Ala Ser Pro Asn Ser Pro Ile Ser Pro Ser Asp Tyr
      50      55      60
Ser Ala Asn Met Asn Val Gly Gly Asp Ser Val Asp Met Leu Leu Ser
65      70      75      80
Ser Val Ser Ala His His Arg Ser Ser Asp Ala Gly Gln Ser Asp Met
      85      90      95
Gly Ser Ile Ser Pro Ser Thr Ala His Thr Thr Pro Asp Ala Thr Thr
      100      105      110
Tyr Lys Thr Ser Asp Glu Glu Asp Ala Thr Gly Lys Ile Thr Thr Pro
      115      120      125
Arg Ser Glu Gly Ser Pro Asn Thr Asn Gly Ser Gly Ser Asp Gly Glu
      130      135      140
Asn Leu Val Cys Lys Trp Gly Pro Cys Gly Lys Thr Phe Gly Ser Ala
145      150      155      160
Glu Lys Leu Tyr Ala His Leu Cys Asp Ala His Val Gly Arg Lys Cys
      165      170      175
Thr His Asn Leu Ser Leu Val Cys Asn Trp Asp Asn Cys Gly Ile Val
      180      185      190
Thr Val Lys Arg Asp His Ile Thr Ser His Ile Arg Val His Val Pro
      195      200      205
Leu Lys Pro Tyr Lys Cys Asp Phe Cys Thr Lys Ser Phe Lys Arg Pro
210      215      220

```

Gln Asp Leu Lys Lys His Val Lys Thr His Ala Asp Asp Asn Glu Gln  
 225 230 235 240  
 Ala His Asn Ala Tyr Ala Lys Pro His Met Gln His Thr His Gln Gln  
 245 250 255  
 Gln Gln Gln Gln Gln Arg Tyr Met Gln Tyr Pro Thr Tyr Ala Ser Gly  
 260 265 270  
 Tyr Glu Tyr Pro Tyr Tyr Arg Tyr Ser Gln Pro Gln Val Gln Val Pro  
 275 280 285  
 Met Val Pro Ser Tyr Ala Ala Val Gly His Met Pro Thr Pro Pro Met  
 290 295 300  
 His Pro His Ala Pro Ile Asp Arg Lys Arg Gln Trp Asp Thr Thr Ser  
 305 310 315 320  
 Asp Phe Phe Asp Asp Ile Lys Arg Ala Arg Val Thr Pro Asn Tyr Ser  
 325 330 335  
 Ser Asp Ile Ala Ser Arg Leu Ser Thr Ile Glu Gln Tyr Ile Gly Ile  
 340 345 350  
 Gln Gly Gln Gln Gln Gln Ala Ser Pro Thr Pro Gln Thr Ala Thr Thr  
 355 360 365  
 Thr Ser Ala Thr Pro Ala Pro Ala Ala Pro His Gln Ala Thr Pro Pro  
 370 375 380  
 Gln Gln Gln Leu Pro Ser Phe Lys Gln Gly Asp Tyr Gln Glu Thr Asp  
 385 390 395 400  
 Gln Phe Leu Asn Gln Leu Gly Ser Asn Ile Tyr Gly Asn Ile Lys Ser  
 405 410 415  
 Val Asp Pro Gln Tyr Glu Ala Pro Ala Glu Phe His Leu Pro His Pro  
 420 425 430  
 Met Gly Tyr Arg Tyr Ala Phe Ser His Ala Pro Ala Pro His Gly Ala  
 435 440 445  
 Ala Pro Val Ala Pro Gln Val Ala Pro Pro Ala His Pro Gly Val His  
 450 455 460  
 Gly Val Ser Ala Pro His Tyr Pro Asp Leu Ser Tyr Ser Arg Ser Thr  
 465 470 475 480  
 Val Pro Gln Leu Ser Ser Arg Phe Glu Asp Val Arg Gln Met Ser Val  
 485 490 495  
 Gly Val Thr Gln Arg Ala Ala Arg Thr Asn Val Glu Glu Ser Asp  
 500 505 510  
 Asp Asp Asp Glu Leu Val Glu Gly Phe Gly Lys Met Ala Ile Ala Asp  
 515 520 525  
 Ser Lys Ala Met Gln Val Ala Gln Met Lys Lys His Leu Glu Val Val  
 530 535 540  
 Ser Tyr Leu Arg Arg Val Leu Gln Glu Ala Arg Glu Thr Glu Ser Gly  
 545 550 555 560  
 Glu Ala Glu Asp Thr Ala Ala Asn Lys Asp Thr Ser Ala Ser Lys Ser  
 565 570 575  
 Ser Leu Tyr Pro Thr Ile Lys Ala Cys  
 580 585

&lt;210&gt; 5

&lt;211&gt; 659

&lt;212&gt; PRT

&lt;213&gt; Candida albicans

&lt;400&gt; 5

Met Asn Tyr Asn Ile His Pro Val Thr Tyr Leu Asn Ala Asp Ser Asn  
 1 5 10 15



```

Thr Gly Ala Ser Glu Ser Thr Ala Ser His His Gly Ser Lys Lys Ser
      20      25      30
Pro Ser Ser Asp Ile Asp Val Asp Asn Ala Xaa Ser Pro Ser Ser Phe
      35      40      45
Thr Ser Ser Gln Ser Pro His Ile Asn Ala Met Gly Asn Ser Pro His
      50      55      60
Ser Ser Phe Thr Ser Gln Ser Ala Ala Asn Ser Pro Ile Thr Asp Ala
      65      70      75      80
Lys Gln His Leu Val Lys Pro Thr Thr Thr Lys Pro Ala Ala Phe Ala
      85      90      95
Pro Ser Ala Asn Gln Ser Asn Thr Thr Ala Pro Gln Ser Tyr Thr Gln
      100      105      110
Pro Ala Gln Gln Leu Pro Thr Gln Leu His Pro Ser Leu Asn Gln Ala
      115      120      125
Tyr Asn Asn Gln Pro Ser Tyr Tyr Leu His Gln Pro Thr Tyr Gly Tyr
      130      135      140
Gln Gln Gln Gln Gln Gln Gln Gln His Gln Glu Phe Asn Gln Pro Ser
      145      150      155      160
Gln Gln Tyr His Asp His His Gly Tyr Tyr Ser Asn Asn Asn Ile Leu
      165      170      175
Asn Gln Asn Gln Pro Ala Pro Gln Gln Asn Pro Val Lys Pro Phe Lys
      180      185      190
Lys Thr Tyr Lys Lys Ile Arg Asp Glu Asp Leu Lys Gly Pro Phe Lys
      195      200      205
Cys Leu Trp Ser Asn Cys Ser Ile Ile Phe Glu Thr Pro Glu Ile Leu
      210      215      220
Tyr Asp His Leu Cys Asp Asp His Val Gly Arg Lys Ser Ser Asn Asn
      225      230      235      240
Leu Ser Leu Thr Cys Leu Trp Glu Asn Cys Gly Thr Thr Thr Val Lys
      245      250      255
Arg Asp His Ile Thr Ser His Leu Arg Val His Val Pro Leu Lys Pro
      260      265      270
Phe His Cys Asp Leu Cys Pro Lys Ser Phe Lys Arg Pro Gln Asp Leu
      275      280      285
Lys Lys His Ser Lys Thr His Ala Glu Asp His Pro Lys Lys Leu Lys
      290      295      300
Lys Ala Gln Arg Glu Leu Met Lys Gln Gln Gln Lys Glu Ala Lys Gln
      305      310      315      320
Gln Gln Lys Leu Ala Asn Lys Arg Ala Asn Ser Met Asn Ala Thr Thr
      325      330      335
Ala Ser Asp Leu Gln Leu Asn Tyr Tyr Ser Gly Asn Pro Ala Asp Gly
      340      345      350
Leu Asn Tyr Asp Asp Thr Ser Lys Lys Arg Arg Tyr Glu Asn Asn Ser
      355      360      365
Gln His Asn Met Tyr Val Val Asn Ser Ile Leu Asn Asp Phe Asn Phe
      370      375      380
Gln Gln Met Ala Gln Ala Pro Gln Gln Pro Gly Val Val Gly Thr Ala
      385      390      395      400
Gly Ser Ala Glu Phe Thr Thr Lys Arg Met Lys Ala Gly Thr Glu Tyr
      405      410      415
Asn Ile Asp Val Phe Asn Lys Leu Asn His Leu Asp Asp His Leu His
      420      425      430
His His His Pro Gln Gln Gln His Pro Gln Gln Gln Tyr Gly Gly Asn
      435      440      445
Ile Tyr Glu Ala Glu Lys Phe Phe Asn Ser Leu Ser Asn Ser Ile Asp

```

```

      450              455              460
Met Gln Tyr Gln Asn Met Ser Thr Gln Tyr Gln Gln His Ala Gly
465              470              475              480
Ser Thr Phe Ala Gln Lys Pro Thr Gln Gln Ala Ser Gly Gln Leu
      485              490              495
Tyr Pro Ser Leu Pro Thr Ile Gly Asn Gly Ser Tyr Thr Ser Gly Ser
      500              505              510
Ser His Lys Glu Gly Leu Val Asn Asn His Asn Gly Tyr Leu Pro Ser
      515              520              525
Tyr Pro Gln Ile Asn Arg Ser Leu Pro Tyr Ser Ser Gly Val Ala Gln
      530              535              540
Gln Pro Pro Ser Ala Leu Glu Phe Gly Gly Val Ser Thr Tyr Gln Lys
545              550              555              560
Ser Ala Gln Ser Tyr Glu Glu Asp Ser Ser Asp Ser Ser Glu Glu Asp
      565              570              575
Asp Tyr Ser Thr Ser Ser Glu Asp Glu Leu Asp Thr Leu Phe Asp Lys
      580              585              590
Leu Asn Ile Asp Asp Asn Lys Val Glu Glu Val Thr Ile Asp Gly Phe
      595              600              605
Asn Leu Lys Asp Val Ala Lys His Arg Glu Met Ile His Ala Val Leu
      610              615              620
Gly Tyr Leu Arg Asn Gln Ile Glu Gln Gln Glu Lys Glu Lys Ser Lys
625              630              635              640
Glu Gln Lys Glu Val Asp Val Asn Glu Thr Lys Leu Tyr Pro Thr Ile
      645              650              655
Thr Ala Phe

```

```

<210> 6
<211> 625
<212> PRT
<213> Saccharomyces cerevisiae

```

```

      <400> 6
Met Val Pro Leu Glu Asp Leu Leu Asn Lys Glu Asn Gly Thr Ala Ala
1      5      10      15
Pro Gln His Ser Arg Glu Ser Ile Val Glu Asn Gly Thr Asp Val Ser
      20      25      30
Asn Val Thr Lys Lys Asp Gly Leu Pro Ser Pro Asn Leu Ser Lys Arg
      35      40      45
Ser Ser Asp Cys Ser Lys Arg Pro Arg Ile Arg Cys Thr Thr Glu Ala
50      55      60
Ile Gly Leu Asn Gly Gln Glu Asp Glu Arg Met Ser Pro Gly Ser Thr
65      70      75      80
Ser Ser Ser Cys Leu Pro Tyr His Ser Thr Ser His Leu Asn Thr Pro
      85      90      95
Pro Tyr Asp Leu Leu Gly Ala Ser Ala Val Ser Pro Thr Thr Ser Ser
      100      105      110
Ser Ser Asp Ser Ser Ser Ser Ser Pro Leu Ala Gln Ala His Asn Pro
      115      120      125
Ala Gly Asp Asp Asp Asp Ala Asp Asn Asp Gly Asp Ser Glu Asp Ile
130      135      140
Thr Leu Tyr Cys Lys Trp Asp Asn Cys Gly Met Ile Phe Asn Gln Pro
145      150      155      160
Glu Leu Leu Tyr Asn His Leu Cys His Asp His Val Gly Arg Lys Ser

```

				165						170					175				
His	Lys	Asn	Leu	Gln	Leu	Asn	Cys	His	Trp	Gly	Asp	Cys	Thr	Thr	Lys				
			180					185					190						
Thr	Glu	Lys	Arg	Asp	His	Ile	Thr	Ser	His	Leu	Arg	Val	His	Val	Pro				
		195					200					205							
Leu	Lys	Pro	Phe	Gly	Cys	Ser	Thr	Cys	Ser	Lys	Lys	Phe	Lys	Arg	Pro				
	210						215					220							
Gln	Asp	Leu	Lys	Lys	His	Leu	Lys	Ile	His	Leu	Glu	Ser	Gly	Gly	Ile				
225					230						235				240				
Leu	Lys	Arg	Lys	Arg	Gly	Pro	Lys	Trp	Gly	Ser	Lys	Arg	Thr	Ser	Lys				
			245						250						255				
Lys	Asn	Lys	Ser	Cys	Ala	Ser	Asp	Ala	Val	Ser	Ser	Cys	Ser	Ala	Ser				
		260						265					270						
Val	Pro	Ser	Ala	Ile	Ala	Gly	Ser	Phe	Lys	Ser	His	Ser	Thr	Ser	Pro				
	275						280					285							
Gln	Ile	Leu	Pro	Pro	Leu	Pro	Val	Gly	Ile	Ser	Gln	His	Leu	Pro	Ser				
	290					295					300								
Gln	Gln	Gln	Gln	Arg	Ala	Ile	Ser	Leu	Asn	Gln	Leu	Cys	Ser	Asp	Glu				
305					310					315					320				
Leu	Ser	Gln	Tyr	Lys	Pro	Val	Tyr	Ser	Pro	Gln	Leu	Ser	Ala	Arg	Leu				
			325					330						335					
Gln	Thr	Ile	Leu	Pro	Pro	Leu	Tyr	Tyr	Asn	Asn	Gly	Ser	Thr	Val	Ser				
		340						345					350						
Gln	Gly	Ala	Asn	Ser	Arg	Ser	Met	Asn	Val	Tyr	Glu	Asp	Gly	Cys	Ser				
	355						360					365							
Asn	Lys	Thr	Ile	Ala	Asn	Ala	Thr	Gln	Phe	Phe	Thr	Lys	Leu	Ser	Arg				
	370					375					380								
Asn	Met	Thr	Asn	Asn	Tyr	Ile	Leu	Gln	Gln	Ser	Gly	Gly	Ser	Thr	Glu				
385					390					395					400				
Ser	Ser	Ser	Ser	Ser	Gly	Arg	Ile	Pro	Val	Ala	Gln	Thr	Ser	Tyr	Val				
			405					410					415						
Gln	Pro	Pro	Asn	Ala	Pro	Ser	Tyr	Gln	Ser	Val	Gln	Gly	Gly	Ser	Ser				
		420						425					430						
Ile	Ser	Ala	Thr	Ala	Asn	Thr	Ala	Thr	Tyr	Val	Pro	Val	Arg	Leu	Ala				
	435						440					445							
Lys	Tyr	Pro	Thr	Gly	Pro	Ser	Leu	Thr	Glu	His	Leu	Pro	Pro	Leu	His				
	450					455					460								
Ser	Asn	Thr	Ala	Gly	Gly	Val	Phe	Asn	Arg	Gln	Ser	Gln	Tyr	Ala	Met				
465					470				475					480					
Pro	His	Tyr	Pro	Ser	Val	Arg	Ala	Ala	Pro	Ser	Tyr	Ser	Ser	Ser	Gly				
			485					490					495						
Cys	Ser	Ile	Leu	Pro	Pro	Leu	Gln	Ser	Lys	Ile	Pro	Met	Leu	Pro	Ser				
		500						505					510						
Arg	Arg	Thr	Met	Ala	Gly	Gly	Thr	Ser	Leu	Lys	Pro	Asn	Trp	Glu	Phe				
	515						520					525							
Ser	Leu	Asn	Gln	Lys	Ser	Cys	Thr	Asn	Asp	Ile	Ile	Met	Ser	Lys	Leu				
	530					535					540								
Ala	Ile	Glu	Glu	Val	Asp	Asp	Glu	Ser	Glu	Ile	Glu	Asp	Asp	Phe	Val				
545					550				555					560					
Glu	Met	Leu	Gly	Ile	Val	Asn	Ile	Ile	Lys	Asp	Tyr	Leu	Leu	Cys	Cys				
			565					570					575						
Val	Met	Glu	Asp	Leu	Asp	Asp	Glu	Glu	Ser	Glu	Asp	Lys	Asp	Glu	Glu				
		580					585					590							
Asn	Ala	Phe	Leu	Gln	Glu	Ser	Leu	Glu	Lys	Leu	Ser	Leu	Gln	Asn	Gln				
	595						600					605							

Met Gly Thr Asn Ser Val Arg Ile Leu Thr Lys Tyr Pro Lys Ile Leu  
 610 615 620  
 Val  
 625

<210> 7  
 <211> 815  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Synthetic primer based on Aspergillus nidulans and  
 herpes virus

<400> 7

Met Ser Ser Arg Gly Ala Met Ala Glu Glu Ala Val Ala Pro Val Ala  
 1 5 10 15  
 Val Pro Thr Thr Gln Glu Gln Pro Thr Ser Gln Pro Ala Ala Ala Gln  
 20 25 30  
 Val Thr Thr Val Thr Ser Pro Ser Val Thr Ala Thr Ala Ala Ala  
 35 40 45  
 Thr Ala Ala Val Ala Ser Pro Gln Ala Asn Gly Asn Ala Ala Ser Pro  
 50 55 60  
 Val Ala Pro Ala Ser Ser Thr Ser Arg Pro Ala Glu Glu Leu Thr Cys  
 65 70 75 80  
 Met Trp Gln Gly Cys Ser Glu Lys Leu Pro Thr Pro Glu Ser Leu Tyr  
 85 90 95  
 Glu His Val Cys Glu Arg His Val Gly Arg Lys Ser Thr Asn Asn Leu  
 100 105 110  
 Asn Leu Thr Cys Gln Trp Gly Ser Cys Arg Thr Thr Thr Val Lys Arg  
 115 120 125  
 Asp His Ile Thr Ser His Ile Arg Val His Val Pro Leu Lys Pro His  
 130 135 140  
 Lys Cys Asp Phe Cys Gly Lys Ala Phe Lys Arg Pro Gln Asp Leu Lys  
 145 150 155 160  
 Lys His Val Lys Thr His Ala Asp Asp Ser Val Leu Val Arg Ser Pro  
 165 170 175  
 Glu Pro Gly Ser Arg Asn Pro Asp Met Phe Gly Gly Asn Gly Lys  
 180 185 190  
 Gly Tyr Ala Ala Ala His Tyr Phe Glu Pro Ala Leu Asn Pro Val Pro  
 195 200 205  
 Ser Gln Gly Tyr Ala His Gly Pro Pro Gln Tyr Tyr Gln Ala His His  
 210 215 220  
 Ala Pro Gln Pro Ser Asn Pro Ser Tyr Gly Asn Val Tyr Tyr Ala Leu  
 225 230 235 240  
 Asn Thr Gly Pro Glu Pro His Gln Ala Ser Tyr Glu Ser Lys Lys Arg  
 245 250 255  
 Gly Tyr Asp Ala Leu Asn Glu Phe Phe Gly Asp Leu Lys Arg Arg Gln  
 260 265 270  
 Phe Asp Pro Asn Ser Tyr Ala Ala Val Gly Gln Arg Leu Leu Ser Leu  
 275 280 285  
 Gln Asn Leu Ser Leu Pro Val Leu Thr Ala Ala Pro Leu Pro Glu Tyr  
 290 295 300  
 Gln Ala Met Pro Ala Pro Val Ala Val Ala Ser Gly Pro Tyr Gly Gly  
 305 310 315 320

Gly Pro His Pro Ala Pro Ala Tyr His Leu Pro Pro Met Ser Asn Val  
 325 330 335  
 Arg Thr Lys Asn Asp Leu Ile Asn Ile Asp Gln Phe Leu Gln Gln Met  
 340 345 350  
 Gln Asp Thr Ile Tyr Glu Asn Asp Asp Asn Val Ala Ala Ala Gly Val  
 355 360 365  
 Ala Gln Pro Gly Ala His Tyr Ile His Asn Gly Ile Ser Tyr Arg Thr  
 370 375 380  
 Thr His Ser Pro Pro Thr Gln Leu Pro Ser Ala His Ala Thr Thr Gln  
 385 390 395 400  
 Thr Thr Ala Gly Pro Ile Ile Ser Asn Thr Ser Ala His Ser Pro Ser  
 405 410 415  
 Ser Ser Thr Pro Ala Leu Thr Pro Pro Ser Ser Ala Gln Ser Tyr Thr  
 420 425 430  
 Ser Gly Arg Ser Pro Ile Ser Leu Pro Ser Ala His Arg Val Ser Pro  
 435 440 445  
 Pro His Glu Ser Gly Ser Ser Met Tyr Pro Arg Leu Pro Ser Ala Thr  
 450 455 460  
 Asp Gly Met Thr Ser Gly Tyr Gly Ser Pro Pro Pro Pro Pro Pro  
 465 470 475 480  
 Pro Pro Pro Pro Ile Lys Val Ala Pro Pro Thr Asp Val Ser Leu Gly  
 485 490 495  
 Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His Ala Asp  
 500 505 510  
 Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp Ser Pro  
 515 520 525  
 Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala Leu Asp  
 530 535 540  
 Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu Gly Ile  
 545 550 555 560  
 Asp Glu Tyr Gly Gly Asp Ile Lys Val Ala Pro Pro Thr Asp Val Ser  
 565 570 575  
 Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His  
 580 585 590  
 Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp  
 595 600 605  
 Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala  
 610 615 620  
 Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu  
 625 630 635 640  
 Gly Ile Asp Glu Tyr Gly Gly Asp Ile Lys Val Ala Pro Pro Thr Asp  
 645 650 655  
 Val Ser Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met  
 660 665 670  
 Ala His Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp  
 675 680 685  
 Gly Asp Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr  
 690 695 700  
 Gly Ala Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp  
 705 710 715 720  
 Ala Leu Gly Ile Asp Glu Tyr Gly Gly Asp Ile Lys Val Ala Pro Pro  
 725 730 735  
 Thr Asp Val Ser Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val  
 740 745 750  
 Ala Met Ala His Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu

	755					760					765								
Gly	Asp	Gly	Asp	Ser	Pro	Gly	Pro	Gly	Phe	Thr	Pro	His	Asp	Ser	Ala				
	770					775					780								
Pro	Tyr	Gly	Ala	Leu	Asp	Met	Ala	Asp	Phe	Glu	Phe	Glu	Gln	Met	Phe				
785					790					795					800				
Thr	Asp	Ala	Leu	Gly	Ile	Asp	Glu	Tyr	Gly	Gly	Asp	Gly	Leu	Gln					
				805					810					815					

<210> 8  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Synthetic primer based on Aspergillus nidulans

<400> 8  
 aactgcagta gttgaccgtg tgattgggtt ct 32

<210> 9  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Synthetic primer based on Aspergillus nidulans

<400> 9  
 ccggaattct ttgtaaactg gcttgaagat 30

<210> 10  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Synthetic oligonucleotide encoding proline rich motif

<400> 10  
 gatccccccc ccctcctcca cccccacccc ctccc 35

<210> 11  
 <211> 31  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Synthetic oligonucleotide encoding proline rich motif

<400> 11  
 gggaggggggt ggggggtggag gagggggggg g 31

<210> 12  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Synthetic primer based on herpes simplex virus  
  
 <400> 12  
 cgcgatatca aagtcgcccc cccgaccgat 30  
  
 <210> 13  
 <211> 30  
 <212> DNA  
 <213> Synthetic primer based on Aspergillus nidulans  
  
 <400> 13  
 cgcgatatcc ccaccgtact cgtcaattcc 30  
  
 <210> 14  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Synthetic primer based on Aspergillus nidulans  
  
 <400> 14  
 tgctctagag gcgccatggc cgaagaagcg 30  
  
 <210> 15  
 <211> 30  
 <212> DNA  
 <213> Aspergillus nidulans  
  
 <400> 15  
 cgcggatccg taaccagaag tcataccgtc 30  
  
 <210> 16  
 <211> 1413  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Synthetic primer based on Aspergillus nidulans  
  
 <400> 16  
 tctagaggcg ccatggccga agaagcggtc gctcctgtag ctgtgcctac gacccaagaa 60  
 caaccaacct ctcaaccgc cgctgcgcag gttacaactg tcacttcgcc ctctgtgact 120  
 gcaacagcgg cggtctgcgac agctgctgtg gccagtcccc aagctaattg caatgctgcc 180  
 tctcctgtcg cccctgcgtc gtcaacatct cgtccagcgg aagaactcac ttgcatgtgg 240  
 caaggctgct ctgagaagct ccctactcca gaatccttat acgaacatgt ctgcgagcgt 300  
 cacgttggcc gaaagagcac gaacaacctc aacctgactt gtcaatgggg tagctgtcgt 360  
 actactactg tgaaacgcga ccatatcacc tctcatatcc ggggtgcacgt tcctctcaag 420  
 ccgcacaagt gtgatttctg tggaaaagcg ttcaagcgtc cccaggattt gaagaagcat 480

```

gttaagacgc acgctgatga ctccggctctg gtacgggtcgc cagagcctgg atctcgcaac      540
ccagatatga tgttcggagg aaatggcaag ggctatgctg ctgcgcacta ttttgagcct      600
gctctcaacc ctgttcccag ccaaggctac gctcatggtc ctccccagta ttaccaggcc      660
catcacgctc cccagccatc gaaccgctct tacggcaacg tctactacgc tctgaatacc      720
ggcccagagc ctcaccaagc gtcgtatgaa tccaagaagc ggggttatga tgcgttaaat      780
gagttctttg gtgacctcaa gcgcgcgacaa tttgacccta attcctacgc tgcgtggggc      840
cagcgcctgc tcagtttgca gaacttgctc ctgcctgttt taacggctgc gcctctgccc      900
gagtaccagg caatgcctgc tcctgtggct gttgctagtg gtccatatgg tggcgccct      960
caccctgcgc cggcatatca tcttccacca atgagcaacg tccgaaccaa gaacgacttg     1020
atcaacatcg accagttcct gcagcaaatg caggacacaa tatatgagaa cgatgataat     1080
gtcgtctcgg ctggtgtcgc tcaacctgga gccattaca ttcataacgg cataagctac     1140
cgcactacac actgcctcc gacacaactt cctcggcac atgccacaac ccagacgact     1200
gctggtccta ttatctcaaa cacatctgcg cactcccctt cgtctagcac tccggctttg     1260
acaccgccct caagtgcgca gtcgtacact tcaggctcgt ctcccatttc acttcctctc     1320
gctcatcgcg tttctccgcc tcatgaaagc ggctccagca tgtaccctcg tctcccttcg     1380
gcgactgacg gtatgacttc tggttacgga tcc                                     1413

```

&lt;210&gt; 17

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Synthetic primer based on *Aspergillus nidulans*

&lt;400&gt; 17

```

ataagaatgc ggccgccctc tgcattattg tctttatc                                     37

```

&lt;210&gt; 18

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Synthetic primer based on *Aspergillus nidulans*

&lt;400&gt; 18

```

tgctctagaa gacattgttg ctatagctgt                                     30

```

&lt;210&gt; 19

&lt;211&gt; 678

&lt;212&gt; DNA

<213> *Aspergillus nidulans*

&lt;400&gt; 19

```

gcggccgccc tctgcattat tgtcttatcc gctattcctg gtgtttttgt tgtcttacta      60
ctttttgtgt cgttgaaatt cttactaggc gttgtgaatc tggatcggat catgctattt     120
tgagggtgtaa tgcattgggtc aaattttctc gagtttcaaa cgaggcagaa gagagatgca     180
gataaatctt gagttttatc atgcagcgaa cgttaccact tatagtttcc ggcagagcac     240
gtaggtcggc ccggcgctcat gtgtagcggg ggagctccag gaccttgagg acgaaaatgg     300
gacggcgatg tataactcca tggaggaacg gacgctgatt ttgtactgtc tgatccgagg     360
ctaattgagaa agcggagggtt caatgtttccc ccggttgatg tcctgaagca gcgaggcccc     420
aagtatcccc tcgtggacat gacatcagtg gtccgactcc cgccgaaccc tctccttca     480
ccggcagccc caccatgtcg ccaaagcaaa tggtagctct gcgattctgg ataccccgcc     540

```



actcaccgtg atacaatttc agcattttgcg aggtgggtctg gtctcctgac gcgctttatt 600  
 tatccctggg ctctcccccac tagctgttcc tgcccgtcca tctctctccg tacagctata 660  
 gcaacaatgt cttctaga 678

<210> 20  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Synthetic primer based on *Aspergillus nidulans*

<400> 20  
 tccccgcgga tggaagcttc gttaaggata att 33

<210> 21  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Synthetic primer based on *Aspergillus nidulans*

<400> 21  
 ataagaatgc ggccgcctac cagattaggg agcatat 37

<210> 22  
 <211> 2229  
 <212> DNA  
 <213> *Aspergillus nidulans*

<400> 22  
 ccgcggatgg aagcttcggt aaggataatt gcctcttttc gaacacctat tcatgttgat 60  
 tagcgatcat tagttatccg gtcgggtaac agaactatgg catactgaac gtcaacttcg 120  
 gaacacgggt ctctcctagt tccggatgga ctaactgccc gtcttccgag aacgtcagct 180  
 atataagtat ctttccccct tcaacgctat cagccatac cttaaagaaa acgcgcagct 240  
 caagcattca gatccacata attagctac tgacgtgaac tatcaaattc catccaccaa 300  
 ttgcccacga tggctcgagat ctccatcccc gcaaaactac ggtacgtccg ccacgggtgtt 360  
 accaaacatt actagccagc tagctcagtc ttaccccggt catgagacca ccccatgcta 420  
 atcatataac gatctttatt atagatatgc catcgccggt tcgctaggcg caatccctgt 480  
 cctgggattc atccatgggt tctcgtcgg ctcttttcgc aaggccgctg gcgtgccgta 540  
 cccccacgcc tatgccagca ttgagcaatg taaagctaac gtgcgtgagc ccaagaaact 600  
 aaatacctat agcaaaacag attgtgttcc aagagagagt actaaatgac gtttgtgaac 660  
 agcccaaagc ctacaaattc aactgcgcac aacgcgcca cggcaacttc ctcgagaacg 720  
 cgccgcagac aatgctctct atcctgggtg caggcgtcaa gtaccagag gcagcagcg 780  
 gcttaggagc ggcctgggtt gttctccgca ccctctacat gctgggctat atttatagcg 840  
 acaagccgaa cggcaccggc aggtacaatg gttcgtgtga cttgcttgcg caagcgggtc 900  
 tttggggatt gagcgcattt ggtgttgcaa aggatttgat gtaaatgtag tgcacatctt 960  
 agcacagagg ggagagttga taaaatgtgg tctgtttgaa tgatagtcgg gttegtgacc 1020  
 tatattcgtg atagtggaga taggtctgcg cctatcttat cgggcccggag caaaaattcc 1080  
 accgcagcgg ggtgagtttt cgttatacag ccataccact tccagcttca aattgtcagt 1140  
 ttaatccagc ccaattcaat cattggagaa ccgccatcat gtcttcgaag tcccacctcc 1200  
 cctacgcaat tcgcgcaacc aaccatccca accctttaac atctaaactc ttctccatcg 1260  
 ccgaggagaa gaaaaccaac gtcaccgtct ccgcagacgt tactacttcc gccgagctcc 1320  
 tcgatcttgc tgaccgtaca tctgcacca atgcccctcc aggataacaa atagctgatg 1380

```

cgtagtgagt acaggcctag gcccctatat cgcagttctg aaaaccacaca tcgacatcct 1440
caccgatctc accccgtcga ccccttcctc gctccaatcc ctccgcgacaa agcacaactt 1500
cctcatcttt gaggaccgca agttcatcga catcggaac accgtgcaaa agcagtacca 1560
cggaggcgct ctccgcatct ccgaatgggc acacatcatc aactgcgcca tcctgccggg 1620
cgaagggatc gtcgaggccc tcgcacagac aaccaagtct cctgacttta aagacgcgaa 1680
tcaacgaggt ctctgattc ttgccgagat gacgagtaag ggatctcttg cgacagggga 1740
gtcacaggca cgctcggttg agtacgcgcg gaagtataag gggtttgtga tgggattcgt 1800
gagtacaagg gcgttgagt aggtgctgcc cgaacagaaa gaggagagcg aggatcttgt 1860
cgtctttacg actggggtga atctgtcgga taagggggat aagctggggc agcagtatca 1920
gacacctggg tcggcggttg ggcgaggtgc ggactttatc attgcgggta ggggcatcta 1980
taaggcggac gatccagtcg aggcggttca gaggtaccgg gaggaaggct ggaaagctta 2040
cgagaaaaga gttggacttt gagtgtgagt ggaaatgtgt aacggtattg actaaaaggg 2100
atccatatgt ttattgcagc cagcatagta ttaccagaaa ggcctcact gacggctcta 2160
gtagtattcg aacagatatt attgtgacca gctctgaacg atatgctccc taatctggta 2220
ggcggccgc 2229

```

<210> 23

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer based on *Aspergillus nidulans*

<400> 23

tgctctagag gcgccatggc cgaagaagcg

30

<210> 24

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer based on *Aspergillus nidulans*

<400> 24

tccccgggg taaccagaag tcataccgtc

30

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/24975**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C07K 14/00; C12N 1/15, 15/63; C12P 21/02

US CL : 435/69.1, 254.11, 320.1; 530/300, 350

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 254.11, 320.1; 530/300, 350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,589,362 A (BUJARD et al) 31 December 1996, see entire document, especially columns 2-14.	1-24
Y	WANG et al. Positive and negative regulation of gene expression in eukaryotic cells with an inducible transcriptional regulator. Gene Therapy. June 1997, Vol. 4, pages 432-441, see entire document, especially pages 432-433.	1-24
Y	GERBER et al. Transcriptional Activation Modulated by Homopolymeric Glutamine and Proline Stretches. Science. 11 February 1994, Vol. 263, pages 808-811, see entire document, especially page 808.	1-24

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  
10 FEBRUARY 1999

Date of mailing of the international search report

25 FEB 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer

TERRY A. MCKELVEY

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/24975

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	TILBURN et al. The Aspergillus PacC zinc finger transcription factor mediates regulation of both acid- and alkaline- expressed genes by ambient pH. The EMBO Journal. 04 February 1995, Vol. 14, No. 4, pages 779-790, see entire document, especially page 779.	1-24
Y	O'REILLY et al. A single serine residue at position 375 of VP16 is critical for complex assembly with Oct-1 and HCF and is a target of phosphorylation by casein kinase II. The EMBO Journal. 01 May 1997, Vol. 16, No. 9, pages 2420-2430, see entire document, especially page 2420.	1, 4, 16, 20
Y	HAO et al. Mutation of Phosphoserine 389 Affects p53 Function in Vivo. The Journal of Biological Chemistry. 15 November 1996, Vol. 271, No. 46, pages 29380-29385, see entire document, especially page 29380.	1, 4, 16, 20
Y	US 5,462,862 A (GROENEN et al) 31 October 1995, see entire document, especially columns 1-3.	9-24

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/24975

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, DIALOG (biotechnology OneSearch databases):

search terms: chimer?, transcription factor?, activat?, pre-activat?, domain?, truncat?, serine, threonine, alanine, aspartic, glutamic, substitut?, replac?, exchang?, PacC, nidulans, fung?, secondary metabolite?, express?, overexpress?, increas?, enhanc?, penicillin, cephalosporin, lovastatin, compactin, cyclosporin, pneumocandin, echinocandin, DNA binding domain?.

